

Rec'd PCT/ATC 28 JAN 2005


Pesticidal Fluoroalkene Derivatives

Description

5

The present invention relates to pesticidal fluoroalkene derivatives of formula I

10



wherein the substituents and the indices have the following meanings:

15

A oxygen or NR^a ;

20

R^a hydrogen; C_1 - C_6 -alkyl, C_2 - C_6 -alkenyl, C_2 - C_6 -alkynyl, wherein the carbon atoms may be partially or fully halogenated;

X hydrogen, halogen; C_1 - C_6 -alkyl or phenyl wherein the alkyl and phenyl groups may be partially or fully halogenated;

25 R^1, R^2 each independently hydrogen, halogen, hydroxyl, cyano, nitro, mercapto, amino; C_1 - C_6 -alkyl, C_2 - C_6 -alkenyl, C_2 - C_6 -alkynyl, C_1 - C_6 -alkoxy, C_2 - C_6 -alkenyloxy, C_1 - C_6 -alkylthio, C_1 - C_6 -alkylamino, di- C_1 - C_6 -alkylamino, C_1 - C_6 -alkoxycarbonyl, C_1 - C_6 -alkylcarbonyloxy, wherein the aliphatic moieties in these 10 substituents are unsubstituted, partially or fully halogenated or substituted by 1 to 3 substituents, each independently selected from R^b ;

30

35 R^b cyano, nitro, halogen, hydroxy, mercapto, amino, carbonyl, aminocarbonyl, aminothiocarbonyl, alkyl, haloalkyl, alkenyl, alkenyloxy, alkynyl, alkoxy, haloalkoxy, alkylthio, alkylamino, dialkylamino, formyl, alkylcarbonyl, alkylsulfonyl, alkoxysulfonyl, alkylsulfonyloxy, alkoxycarbonyl, alkylcarbonyloxy, alkylaminocarbonyl, dialkylaminocarbonyl, alkylaminothiocarbonyl, dialkylaminothiocarbonyl, alkylenedioxy or cycloalkyl, wherein the alkyl groups in these radicals contain 1 to 6 carbon atoms and the abovementioned alkenyl or alkynyl groups in these radicals contain 2 to 6 carbon atoms, and wherein the carbon atoms in these groups may be partially or fully halogenated;

40

45

2

Het a monocyclic or bicyclic 3- to 10-membered heteroaromatic ring system containing 1 to 5 heteroatoms selected from oxygen, sulfur and nitrogen, unsubstituted, partially or fully halogenated or substituted by 1 to 4 substituents, each independently selected from R^c:

R^c R^b, C₁-C₆-alkoxy-C₁-C₆-alkyl, C₁-C₆-alkylsulfinyl, C₁-C₆-alkylaminosulfonyl, di-C₁-C₆-alkylaminosulfonyl, C₁-C₆-alkylcarbonylamino, wherein the last mentioned 5 carbon chains and those defined under R^b are unsubstituted, partially or fully halogenated or substituted by from 1 to 3 cyano, hydroxy, mercapto, amino, C₁-C₆-alkylthio, C₁-C₆-alkylamino, di-C₁-C₆-alkylamino, C₁-C₆-alkoxycarbonyl, C₁-C₆-alkylcarbonyloxy or nitro groups;

cycloalkyl, cycloalkoxy, saturated or partially unsaturated heterocyclyl, heterocyclyloxy, wherein the cyclic systems contain 3 to 10 ring members, and the carbon atoms in the heterocycles may be substituted by 1 to 4 heteroatoms selected from nitrogen, sulfur and oxygen,

aryl, aryloxy, arylthio, aryl-C₁-C₆-alkoxy, aryl-C₁-C₆-alkyl, wherein the mono- or bicyclic ring systems contain 5 to 10 ring members,

hetaryl, hetaryloxy, hetarylthio, wherein the mono- or bicyclic ring systems contain 5 to 10 ring members wherein 1 to 3 carbon atoms may be substituted by heteroatoms selected from nitrogen, sulfur and oxygen,

and wherein the cyclic, aromatic and heteroaromatic systems may be partially or fully halogenated or may be substituted by from 1 to 3 groups selected from halogen, cyano, nitro, hydroxy; C₁-C₆-alkyl, C₁-C₆-alkoxy, C₁-C₆-alkylthio, C₁-C₆-alkylamino, C₁-C₆-alkylcarbonyl, C₁-C₆-alkoxycarbonyl, di-C₁-C₆-alkylamino, C₂-C₆-alkenyl, C₂-C₆-alkenyloxy and C₂-C₆-alkynyl, wherein the carbon atoms of these substituents may be partially or fully halogenated;

m 0, 1 or 2;

n 0, 1, 2, or 3;

p 0, 1, 2, 3, 4, 5, or 6.

Furthermore, the present invention relates to processes for the preparation of compounds of formula I, compositions containing them and their use for the control of pests such as nematodes,

3

insects, arachnids, harmful fungi and unwanted plants, and the protection of plants from those pests as well as their use for treating, controlling, preventing and protecting warm-blooded animals, fish and humans against infestation and infection by helminths, arachnids and arthropod endo- and ectoparasites.

In WO-A 86/07590, nematicidal halosubstituted alkene derivatives are disclosed, some of which are substituted by a carbonyloxy group substituted by dihydrothiazolylthiomethylene.

10

The compounds of formula I differ from the compounds known from WO-A 86/07590 in that they are substituted by aromatic heterocyclic substituents.

15 In WO 97/08130, pesticidal 4,4-difluoro-3-butenylester derivatives and 4,4-difluoro-3-halogen-3-butenylester derivatives are described some of which carry a hetarylthioalkyl substituent on the carbonyl group.

20 Contrary to the compounds disclosed in WO-A 97/08130, in compounds of formula I a hetaryl moiety is bonded to the carbonylalkyl backbone via a sulfur atom.

Furthermore, EP-A 1000946 teaches 2-(substituted thio)thiazolo-[4,5-b]pyridine compounds which may bear a haloalkenyl-oxycarbonyl-alkylthio radical as the substituting thio group.


However, the pesticidal activity of the compounds known from the above literature in many cases is unsatisfactory.

It is an object of the present invention to provide compounds having improved nematicidal, insecticidal and acaricidal activity. It is also an object to provide compounds for controlling harmful fungi, unwanted plants and parasites.

We have found that this object is achieved by the fluoroalkene derivatives of formula I. Furthermore, we have found processes for preparing the compounds of formula I and the use of the compounds I and compositions comprising them for use for the control of nematodes, insects, arachnids, harmful fungi and unwanted plants and the protection of plants from those pests as well as for treating, controlling, preventing and protecting warm-blooded animals, fish and humans against infestation and infection by helminths, acarids and arthropod endo- and ectoparasites.

4


Fluoroalkene derivatives of formula I.1,




wherein A, X, R¹, R², Het, n and p are as defined for formula I above, are obtainable by, in a first step, reaction of compounds of formula II



wherein Het is as defined for formula I, with compounds of formula III



wherein R¹, R² and n are as defined for formula I and L is a nucleophilic exchangeable leaving group, preferably halogen such as bromine, and Rⁱ is hydrogen, C₁-C₆-alkyl or aryl-C₁-C₆-alkyl, such as benzyl, in the presence of a base to give sulfide compounds of formula IV.



The reaction to sulfides IV is usually carried out at temperatures of from 0°C to 150°C, preferably from 15°C to 80°C, in an inert organic solvent in the presence of a base.

Suitable solvents are halogenated hydrocarbons, such as methylene chloride and chlorobenzene, ethers, such as dimethylether, diglyme, dioxane and tetrahydrofuran, nitriles, such as acetonitrile, ketones, such as acetone, and also dimethyl sulfoxide, dimethyl formamide and dimethyl acetamide. Preferred solvents are acetone and dimethyl formamide. It is also possible to use mixtures of the solvents mentioned.

Suitable bases are, inorganic compounds, such as alkali metal and alkaline earth metal carbonates, such as lithium carbonate, potassium carbonate and calcium carbonate, alkali metal bicarbonates, such as sodium bicarbonate, and also organic bases, such as tertiary amines, such as trimethyl amine, triethyl amine, tri-isopropyl ethyl amine und N-methyl- piperidine, and pyridine. Particular preference is given to alkaline earth metal carbonates, especially potassium carbonate.

5

In general, the base is employed in equimolar amounts or in excess.

The starting materials are generally reacted with one another in 5 equimolar amounts. In terms of yield, it may be advantageous to use an excess of compounds of formula III based on compounds Het-SH.

Heterocyclic thiols of formula II are known, or are commercially 10 available, or they can be prepared by known methods [see e.g. Synthesis 3, 358-360 (2001)].

Compounds of formula IV are known from the literature or are commercially available [see e.g. J. Org. Chem. 62, 9173-9176 15 (1997)].

Compounds of formula IV wherein Rⁱ is C₁-C₆-alkyl or aryl-C₁-C₆-alkyl are hydrolyzed to compounds IV wherein Rⁱ is H, for example in the presence of an aqueous acid or base, and reacted 20 with fluoroalkenyl compounds of formula V,



25 wherein X and p are as defined for formula I and Y is (a) a nucleophilically exchangeable group, such as halogen or hydroxy, to yield compounds I wherein A is oxygen, or (b) an amino group NHR^a, wherein R^a is as defined for formula I, preferably hydrogen, to yield compounds I wherein A is NR^a.

30

The reaction is carried out by common coupling methods such as in the presence of a base, optionally under activating conditions, such as by converting carboxylic acids of formula V into their corresponding carboxylic acid halides or by means of dehydrating 35 agents such as carbodiimides, to give compounds of formula Ia [lit.: J. March, Advanced Organic Chemistry: reactions, mechanisms and structure, 4th ed. 1992, Wiley&Sons, New York].

The reaction is usually carried out at temperatures of from 0°C to 40 150°C, preferably from 20°C to 60°C, in an inert organic solvent in the presence of a base.

Suitable solvents are nitriles, such as acetonitrile and propionitrile, and also dimethyl sulfoxide, dimethyl formamide 45 and dimethyl acetamide. Preferred solvents are dimethylether and

6

acetonitrile. It is also possible to use mixtures of the solvents mentioned.

The starting materials are generally reacted with one another in 5 equimolar amounts. In terms of yield, it may be advantageous to use an excess of compounds V, based on acids IV.

Compounds of formula V are known from the literature or are commercially available [see e.g. WO 86/07590 and WO 95/24403].

- 10 Sulfinyl and sulfonyl compounds of formula I wherein m is 1 or 2 may be prepared by oxidizing compounds of formula Ia. The oxidation is usually carried out at temperatures of from -10°C to 150°C, preferably from 0°C to 60°C, in an inert organic solvent or water. Suitable oxidizing agents are, for example m-chloroperbenzoic acid, peracetic acid, $\text{H}_2\text{O}_2 \times \text{BF}_3$, $\text{K}_2\text{S}_2\text{O}_7/\text{H}_2\text{SO}_4$, peroxytrifluoroacetic acid, or hydrogen peroxide, optionally in combination with catalytic amounts of sodium tungsten dihydrate.

- Suitable solvents are halogenated hydrocarbons, such as methylene 20 chloride and chloroform alcohols, such as methanol and tert.-butanol, carboxylic acids such as acetic acid and trifluoroacetic acid, and also dimethyl sulfoxide, dimethyl formamide and dimethyl acetamide. Preferred solvents are methylene chloride and acetic acid. It is also possible to use 25 mixtures of the solvents mentioned.

The reaction mixtures are worked up in a customary manner, for example by mixing with water, phase separation and, if appropriate, chromatographic purification of the crude products.

- 30 In some cases, the intermediates and end products are obtained in the form of colorless or pale brown viscous oils, which are purified or freed from volatile components under reduced pressure and at moderately elevated temperature. If the intermediates and end products are obtained as solids, they can also be purified by 35 recrystallization or digestion.

If individual compounds I are not obtainable by the routes described above, they can be prepared by derivatization of other compounds I.

40

In the definitions of the symbols given in the above formulae, collective terms were used which generally represent the following substituents:

- 45 **Halogen:** fluorine, chlorine, bromine and iodine;

7

Alkyl: saturated, straight-chain or branched hydrocarbon radicals having 1 to 4 or 6 carbon atoms, such as methyl, ethyl, propyl, 1-methylethyl, butyl, 1-methylpropyl, 2-methylpropyl, 1,1-dimethylethyl, pentyl, 1-methylbutyl, 2-methylbutyl, 5 3-methylbutyl, 2,2-dimethylpropyl, 1-ethylpropyl, hexyl, 1,1-dimethylpropyl, 1,2-dimethylpropyl, 1-methylpentyl, 2-methylpentyl, 3-methylpentyl, 4-methylpentyl, 1,1-dimethylbutyl, 1,2-dimethylbutyl, 1,3-dimethylbutyl, 2,2-dimethylbutyl, 2,3-dimethylbutyl, 3,3-dimethylbutyl, 10 1-ethylbutyl, 2-ethylbutyl, 1,1,2-trimethylpropyl, 1,2,2-trimethylpropyl, 1-ethyl-1-methylpropyl and 1-ethyl-2-methylpropyl;

Haloalkyl: straight-chain or branched alkyl groups having 1 to 4 15 or 6 carbon atoms (as mentioned above), where some or all of the hydrogen atoms in these groups may be replaced by halogen atoms as mentioned above, for example C₁-C₂-haloalkyl, such as chloromethyl, bromomethyl, dichloromethyl, trichloromethyl, fluoromethyl, difluoromethyl, trifluoromethyl, 20 chlorofluoromethyl, dichlorofluoromethyl, chlorodifluoromethyl, 1-chloroethyl, 1-bromoethyl, 1-fluoroethyl, 2-fluoroethyl, 2,2-difluoroethyl, 2,2,2-trifluoroethyl, 2-chloro-2-fluoroethyl, 2-chloro-2,2-difluoroethyl, 2,2-dichloro-2-fluoroethyl, 2,2,2-trichloroethyl and pentafluoroethyl;

25

Alkenyl: unsaturated, straight-chain or branched hydrocarbon radicals having 2 to 6 carbon atoms and a double bond in any position, such as ethenyl, 1-propenyl, 2-propenyl, 1-methyl-ethenyl, 1-butenyl, 2-butenyl, 3-butenyl, 1-methyl-1-propenyl, 30 2-methyl-1-propenyl, 1-methyl-2-propenyl, 2-methyl-2-propenyl; 1-pentenyl, 2-pentenyl, 3-pentenyl, 4-pentenyl, 1-methyl-1-butenyl, 2-methyl-1-butenyl, 3-methyl-1-butenyl, 1-methyl-2-butenyl, 2-methyl-2-butenyl, 3-methyl-2-butenyl, 1-methyl-3-butenyl, 2-methyl-3-butenyl, 3-methyl-3-butenyl, 1,1-dimethyl-35 2-propenyl, 1,2-dimethyl-1-propenyl, 1,2-dimethyl-2-propenyl, 1-ethyl-1-propenyl, 1-ethyl-2-propenyl, 1-hexenyl, 2-hexenyl, 3-hexenyl, 4-hexenyl, 5-hexenyl, 1-methyl-1-pentenyl, 2-methyl-1-pentenyl, 3-methyl-1-pentenyl, 4-methyl-1-pentenyl, 1-methyl-2-pentenyl, 2-methyl-2-pentenyl, 3-methyl-2-pentenyl, 4-methyl-40 2-pentenyl, 1-methyl-3-pentenyl, 2-methyl-3-pentenyl, 3-methyl-3-pentenyl, 4-methyl-3-pentenyl, 1-methyl-4-pentenyl, 2-methyl-4-pentenyl, 3-methyl-4-pentenyl, 4-methyl-4-pentenyl, 1,1-dimethyl-2-butenyl, 1,1-dimethyl-3-butenyl, 1,2-dimethyl-1-butenyl, 1,2-dimethyl-2-butenyl, 1,2-dimethyl-3-butenyl, 45 1,3-dimethyl-1-butenyl, 1,3-dimethyl-2-butenyl, 1,3-dimethyl-3-butenyl, 2,2-dimethyl-3-butenyl, 2,3-dimethyl-1-butenyl, 2,3-dimethyl-2-butenyl, 2,3-dimethyl-3-butenyl, 3,3-dimethyl-

8

1-butenyl, 3,3-dimethyl-2-butenyl, 1-ethyl-1-butenyl, 1-ethyl-2-butenyl, 1-ethyl-3-butenyl, 2-ethyl-1-butenyl, 2-ethyl-2-butenyl, 2-ethyl-3-butenyl, 1,1,2-trimethyl-2-propenyl, 1-ethyl-1-methyl-2-propenyl, 1-ethyl-2-methyl-1-propenyl and
 5 1-ethyl-2-methyl-2-propenyl;

Haloalkenyl: unsaturated, straight-chain or branched hydrocarbon radicals having 2 to 6 carbon atoms and a double bond in any position (as mentioned above), where some or all of the hydrogen
 10 atoms in these groups may be replaced by halogen atoms as mentioned above, in particular by fluorine, chlorine and bromine;

Alkynyl: straight-chain or branched hydrocarbon groups having 2 to 6 carbon atoms and a triple bond in any position, such as
 15 ethynyl, 1-propynyl, 2-propynyl, 1-butyne, 2-butyne, 3-butyne, 1-methyl-2-propynyl, 1-pentyne, 2-pentyne, 3-pentyne, 4-pentyne, 1-methyl-2-butyne, 1-methyl-3-butyne, 2-methyl-3-butyne, 3-methyl-1-butyne, 1,1-dimethyl-2-propynyl, 1-ethyl-2-propynyl, 1-hexynyl, 2-hexynyl, 3-hexynyl, 4-hexynyl, 5-hexynyl,
 20 1-methyl-2-pentyne, 1-methyl-3-pentyne, 1-methyl-4-pentyne, 2-methyl-3-pentyne, 2-methyl-4-pentyne, 3-methyl-1-pentyne, 3-methyl-4-pentyne, 4-methyl-1-pentyne, 4-methyl-2-pentyne, 1,1-dimethyl-2-butyne, 1,1-dimethyl-3-butyne, 1,2-dimethyl-3-butyne, 2,2-dimethyl-3-butyne, 3,3-dimethyl-1-butyne,
 25 1-ethyl-2-butyne, 1-ethyl-3-butyne, 2-ethyl-3-butyne and 1-ethyl-1-methyl-2-propynyl;

Haloalkynyl: straight-chain or branched hydrocarbon groups having 2 to 6 carbon atoms and a triple bond in any position (as
 30 mentioned above), where some or all of the hydrogen atoms in these groups may be replaced by halogen atoms as mentioned above, in particular by fluorine, chlorine and bromine;

Alkoxy carbonyl: straight-chain or branched alkoxy groups having 1
 35 to 6 carbon atoms (as mentioned above) which are attached to the skeleton via a carbonyl group (-CO-);

Alkylaminocarbonyl: straight-chain or branched alkylamino groups having 1 to 6 carbon atoms (as mentioned above) which are
 40 attached to the skeleton via a carbonyl group (-CO-);

Alkylcarbonyloxy: straight-chain or branched alkyl groups having 1 to 6 carbon atoms (as mentioned above) which are attached to the skeleton via a carbonyloxy group (-C(=O)O-);

Cycloalkyl: monocyclic or bicyclic saturated hydrocarbon groups having 3 to 6, 8 or 10 carbon ring atoms, e.g. C₃-C₈-cycloalkyl such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl and dihydronaphthalin;

5

Heterocyclyl: 5- to 10-membered saturated or partially unsaturated rings which besides carbon ring atoms contain from 1 to 3 or 4 heteroatoms selected from nitrogen, oxygen and sulfur, e.g. from 1 to 3 nitrogen atoms and/or 1 oxygen or sulfur atom
10 and/or 1 or 2 oxygen and/or sulfur atoms;

Aryl: mono- or bicyclic aromatic ringsystems containing 5 to 10 carbon ring atoms, e.g. phenyl or naphthyl;

15 **Arylmethylene:** mono- or bicyclic aromatic ringsystems containing 5 to 14 carbon ring atoms (as mentioned above) which are attached to the skeleton via a methylene (-CH₂-) group.

Hetaryl: a monocyclic or bicyclic 3- to 10-membered
20 heteroaromatic ring system containing 1 to 5 heteroatoms selected from oxygen, sulfur and nitrogen,
e.g. 5-membered hetaryl containing beside carbon atoms 1 to 3 nitrogen atoms and/or 1 sulfur- or oxygen atom, wherein the ring system may be bonded to the backbone via carbon or nitrogen, such
25 as furyl, thienyl, pyrrolyl, isoxazolyl, isothiazolyl, pyrazolyl, oxazolyl, thiazolyl, imidazolyl, 1,2,4-oxadiazolyl, 1,2,4-thiadiazolyl, 1,2,4-triazolyl, 1,3,4-oxadiazolyl, 1,3,4-thiadiazolyl and 1,3,4-triazolyl; or
e.g. 5-membered hetaryl containing beside carbon atoms 1 to 3
30 nitrogen atoms and/or 1 sulfur- or oxygen atom, wherein the ring system may be bonded to the backbone via carbon or nitrogen, wherein 2 adjacent ring members are bridged by a buta-1,3-dien-1,4-diyl group, wherein 1 or 2 carbon atoms may be substituted by nitrogen atoms, such as benzofuranyl,
35 isobenzofuranyl, benzothienyl, isobenzothienyl, indolyl, isoindolyl, indazolyl, indoleninyl, isobenzazolyl, pyranopyrrolyl, isoindazolyl, indoxazinyl, benzoxazolyl, benzothiazolyl, benzimidazolyl, pyridoxazolyl, pyridothiazolyl, pyrazinoxazolyl, pyrazinthiazolyl, pyridazinoxazolyl,
40 pyridazinthiazolyl, pyrimidinoxazolyl, pyrimidinthiazolyl, pyrimidinazolyl, benzopyranyl, purinyl;

Alkylsulfinyl: straight-chain or branched alkyl groups having 1 to 6 carbon atoms (as mentioned above) which are attached to the
45 skeleton via a sulfinyl group (-SO-);

10

Alkylsulfonyl: straight-chain or branched alkyl groups having 1 to 6 carbon atoms (as mentioned above) which are attached to the skeleton via a sulfonyl group ($-\text{SO}_2-$);

5 Alkylsulfonyloxy: straight-chain or branched alkylsulfonyl groups having 1 to 6 carbon atoms (as mentioned above) which are attached to the skeleton via an oxygen atom;

Alkoxy sulfonyl: straight-chain or branched alkoxy groups having 1 to 6 carbon atoms (as mentioned above) which are attached to the skeleton via a sulfonyl group ($-\text{SO}_2-$);

15 Alkylaminosulfonyl: straight-chain or branched alkylamino groups having 1 to 6 carbon atoms (as mentioned above) which are attached to the skeleton via a sulfonyl group ($-\text{SO}_2-$);

With respect to the intended use of the fluoroalkene derivatives of formula I, particular preference is given to the following meanings of the substituents, in each case on their own or in combination:

Preference is given to compounds of formula I wherein A is oxygen or NH.

25 Particular preference is given to compounds of formula I wherein A is oxygen.

Preference is given to compounds of the formula I in which R^a is hydrogen or C_1 - C_6 -alkyl, hydrogen being preferred most.

30

Furthermore, preference is given to compounds of formula I wherein X is hydrogen, halogen, C_1 - C_4 -alkyl, C_1 - C_4 -haloalkyl, or phenyl.

35 Particular preference is given to compounds of formula I wherein X is hydrogen or halogen, especially fluorine.

Especially preferred are compounds of formula I wherein X is fluorine.

40

Preference is given to compounds of formula I wherein R^1 and R^2 are each independently hydrogen, halogen, C_1 - C_6 -alkyl, or phenyl wherein the alkyl and phenyl group are unsubstituted, partially or fully halogenated.

45

11

Particular preference is given to compounds of formula I wherein R¹ and R² are each independently hydrogen, halogen, C₁-C₄-alkyl, or phenyl.

- 5 Especially preferred are compounds of formula I wherein R¹ and R² are each independently hydrogen or C₁-C₄-alkyl.

Particular preference is given to compounds of formula I wherein one of R¹ and R² is not hydrogen.

10

Particular preference is given to compounds of the formula I wherein R^b is halogen, hydroxy, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkoxy or C₁-C₆-haloalkoxy.

- 15 Especially preferred are compounds of the formula I wherein R¹ and R² are substituted by hydrogen or C₁-C₆-alkyl, hydrogen being preferred most.

Preference is given to compounds of the formula I wherein Het is
20 bonded via carbon.

- Particular preference is given to compounds of the formula I wherein Het is oxazolyl, thiazolyl, imidazolyl, pyrimidinyl, pyridyl, purinyl, pyridazinonyl, benzoxazolyl, pyridoxazolyl,
25 benzothiazolyl, pyridothiazolyl, benzimidazolyl, pyrimidinazolyl, 1,3,4-oxadiazol-2-yl, or 1,3,4-thiadiazol-2-yl, each ring system being unsubstituted, partially or fully halogenated or substituted by from 1 to 3 or 4 groups selected from R^c as defined for formula I above.

30

- Particular preference is given to compounds of the formula I wherein Het is 2-oxazolyl, 2-thiazolyl, 2-imidazolyl, 2-pyrimidinyl, 2-pyridyl, 2-purinyl, 5-pyridazinonyl, 2-benzoxazolyl, 2-pyridoxazolyl, 2-benzothiazolyl,
35 2-benzimidazolyl, 2-pyrimidinazolyl, 1,3,4-oxadiazol-2-yl, or 1,3,4-thiadiazol-2-yl, each ring system being unsubstituted, partially or fully halogenated or substituted by from 1 to 3 or 4 groups selected from R^c as defined for formula I above.

- 40 Especially preferred are compounds of formula I wherein Het is 2-oxazolyl, 2-thiazolyl, 2-benzoxazolyl, 2-pyridoxazolyl, 2-benzothiazolyl, 1,3,4-oxadiazol-2-yl, or 1,3,4-thiadiazol-2-yl, each ring system being unsubstituted, partially or fully halogenated or substituted by from 1 to 2 or 3 groups selected
45 from R^c as defined for formula I above.

12

Preference is given to compounds of the formula I wherein R^c is cyano, nitro, halogen, hydroxy, mercapto, amino, carboxyl, aminocarbonyl, alkyl, haloalkyl, alkoxyalkyl, alkenyl, alkenyloxy, alkynyl, alkoxy, haloalkoxy, alkylthio, alkylamino, dialkylamino, 5 formyl, alkylcarbonyl, alkylsulfonyl, alkoxycarbonyl, alkylcarbonyloxy, alkylaminocarbonyl, or dialkylaminocarbonyl, wherein the alkyl groups in these radicals contain 1 to 6 carbon atoms and the abovementioned alkenyl or alkynyl groups in these radicals contain 2 to 6 carbon atoms, and wherein the carbon 10 atoms in these groups may be partially or fully halogenated, or 5- to 10-membered mono- or bicyclic aryl, or 5- to 10-membered mono- or bicyclic hetaryl, wherein 1 to 3 carbon atoms may be substituted by heteroatoms selected from nitrogen, sulfur and oxygen, wherein the aryl or hetaryl ring systems may be partially 15 or fully halogenated or may be substituted by from 1 to 3 groups selected from halogen, cyano, nitro, hydroxy, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkoxy, or C₁-C₆-haloalkoxy.

Particular preference is given to compounds of the formula I 20 wherein R^c is hydroxy, halogen, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkoxy, C₁-C₆-haloalkoxy and phenyl, unsubstituted, partially or fully halogenated or substituted by from 1 to 3 nitro, hydroxy, cyano, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkoxy, C₁-C₆-haloalkoxy, C₁-C₆-alkylamino, or di-C₁-C₆-alkylamino.

25 Particular preference is given to compounds of the formula I in which m is 0 or 2.

Particular preference is given to compounds of the formula I in 30 which m is zero.

Preference is given to compounds of the formula I in which n is 0 or 1.

35 Particular preference is given to compounds of the formula I in which n is zero.

Preference is given to compounds of the formula I in which p is 2 or 4.

40 Preference is given to compounds of the formula I in which p is 2.

Moreover, fluoroalkene derivatives of formula I are preferred 45 wherein the substituents and the indices have the following meanings:

A oxygen or NH;

13

X hydrogen or halogen;

R¹, R² each independently hydrogen, C₁-C₆-alkyl, or phenyl which is unsubstituted or substituted by one to three halogen atoms;

Het oxazolyl, thiazolyl, imidazolyl, pyrimidinyl, pyridyl, purinyl, pyridazinonyl, benzoxazolyl, pyridoxazolyl, benzothiazolyl, pyridothiazolyl, benzimidazolyl, pyrimidazolyl, oxadiazolyl, or thiadiazolyl, each ring being unsubstituted, partially or fully halogenated or substituted by from one to three groups selected from R^c.

10 R^c is hydroxy, halogen, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkoxy, C₁-C₆-haloalkoxy and phenyl, unsubstituted, partially or fully halogenated or substituted by from 1 to 3 nitro, hydroxy, cyano, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkoxy, C₁-C₆-haloalkoxy, C₁-C₆-alkylamino, or di-C₁-C₆-alkylamino.

m, n 0, 1 or 2;

p 2 or 4;

Likewise, particular preference is given to compounds of the
20 formula I wherein substituents and the indices have the following meanings:

A oxygen;

X hydrogen or fluorine;

R¹, R² each independently hydrogen, C₁-C₄-alkyl;

25 Het 2-oxazolyl, 2-thiazolyl, 2-benzoxazolyl, 2-pyridoxazolyl, 2-benzothiazolyl, 2-pyridothiazolyl, 1,3,4-oxadiazol-2-yl, or 1,3,4-thiadiazol-2-yl, each ring being unsubstituted, partially or fully halogenated or substituted by from one to three groups selected from hydroxy, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkoxy, C₁-C₆-haloalkoxy, or phenyl, unsubstituted,
30 partially or fully halogenated or substituted by from 1 to 3 nitro, cyano, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkoxy or C₁-C₆-haloalkoxy groups.

m, n zero;

35 p 2 or 4.

Particular preference is given to compounds of formula I wherein A is oxygen, X is hydrogen or fluorine, R¹ and R² are each independently hydrogen, C₁-C₄-alkyl, or phenyl, Het is 2-benzoxazolyl,
40 2-benzothiazolyl, 4-pyridothiazol-2-yl, 4-pyridoxazol-2-yl, 5-pyridothiazol-2-yl, 5-pyridoxazol-2-yl, 6-pyridothiazol-2-yl, 6-pyridoxazol-2-yl, 7-pyridothiazol-2-yl, 7-pyridoxazol-2-yl, 2-pyrazinthiazolyl, 2-pyrazinoxazolyl, each ring being unsubstituted or substituted by 1 group selected from halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-haloalkoxy, C₁-C₄-alkoxy, nitro, amino,
45 and methylcarbonylamino, m is zero, n is 1, and p is 2 or 4.

14


Particular preference is given to compounds of formula I wherein A is oxygen, X is hydrogen or fluorine, R¹ and R² are each independently hydrogen, C₁-C₄-alkyl, or phenyl, Het is 2-oxazolyl, 2-thiazolyl, 2-imidazolyl, each ring being unsubstituted or substituted by 1 or 2 groups selected from halogen, C₁-C₄-alkyl, C₁-C₄-alkoxymethylene, C₁-C₄-alkylcarbonyloxy, and phenyl which may be substituted by from 1 to 3 substituents selected from halogen, C₁-C₄-alkyl and hydroxy, m is zero, n is 1, and p is 2 or 4.

10

Particular preference is given to compounds of formula I wherein A is oxygen, X is hydrogen or fluorine, R¹ and R² are each independently hydrogen, C₁-C₄-alkyl, or phenyl, Het is thiadiazolyl or oxadiazolyl, each ring being unsubstituted or substituted by one group selected from halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-haloalkoxy, C₁-C₄-alkoxy, nitro, amino, and methylcarbonylamino, m and n are integers of 0, and p is an integer of 2 or 4.


Furthermore, compounds of the following formulae are especially preferred:

Compounds of formula I.2



wherein B is nitrogen or CH, in particular CH, D is nitrogen or CH, in particular CH, E is oxygen or sulfur, R¹ and R² are each independently hydrogen, C₁-C₄-alkyl or phenyl, q is 0 or 1, R^c is halogen, methyl, methoxy, ethoxy, trifluoromethyl, trifluoromethoxy, m is 0, 1 or 2, n is 0 or 1 and p is 2 or 4, with the proviso that B is CH when E is sulfur.


35 Compounds of formula I.3



wherein E' is oxygen, sulfur or NH, R¹ and R² are each independently hydrogen or C₁-C₄-alkyl, q is 0, 1, or 2, R^c is phenyl which may be substituted by one or two halogen, m is 0, 1 or 2, n is 0 or 1, p is 2 or 4.

15

Compounds of formula I.4



5

wherein E is oxygen or sulfur, R^1 and R^2 are each independently hydrogen or C_1 - C_4 -alkyl, R^c is C_1 - C_4 -alkyl or phenyl which may be substituted by halogen or methyl, and p is 2.

10 With respect to their use, particular preference is given to the compounds I.5, I.6 and I.7 compiled in the Tables below.


Moreover, the groups mentioned for a substituent in the Tables are on their own, independently of the combination in which they are mentioned, a particularly preferred embodiment of the

15 substituent in question.

Further preference is given to the compounds of the tables 1 to 440 wherein A is NH.

20 Table 1

Compounds of the formula I.5



25

wherein X is hydrogen, p is 2, R^1 and R^2 are hydrogen and the combination of B, D, E and R^c for a compound corresponds in each case to a row of Table A.

30

Table 2

Compounds of the formula I.5 wherein X is hydrogen, p is 2, R^1 is hydrogen and R^2 is methyl and the combination of B, D, E and R^c for a compound corresponds in each case to a row of Table A.

35

Table 3

Compounds of the formula I.5 wherein X is hydrogen, p is 2, R^1 is hydrogen and R^2 is ethyl and the combination of B, D, E and R^c for a compound corresponds in each case to a row of Table A.

40

Table 4

Compounds of the formula I.5 wherein X is hydrogen, p is 2, R^1 is hydrogen and R^2 is i-propyl and the combination of B, D, E and R^c for a compound corresponds in each case to a row of Table A.

45

Table 5

Compounds of the formula I.5 wherein X is hydrogen, p is 2, R^1 is

16

hydrogen and R^2 is n-propyl and the combination of B, D, E and R^c for a compound corresponds in each case to a row of Table A.

Table 6

- 5 Compounds of the formula I.5 wherein X is hydrogen, p is 2, R^1 is hydrogen and R^2 is cyclo-propyl and the combination of B, D, E and R^c for a compound corresponds in each case to a row of Table A.

Table 7

- 10 Compounds of the formula I.5 wherein X is hydrogen, p is 2, R^1 is hydrogen and R^2 is n-butyl and the combination of B, D, E and R^c for a compound corresponds in each case to a row of Table A.

Table 8

- 15 Compounds of the formula I.5 wherein X is hydrogen, p is 2, R^1 is hydrogen and R^2 is i-butyl and the combination of B, D, E and R^c for a compound corresponds in each case to a row of Table A.

Table 9

- 20 Compounds of the formula I.5 wherein X is hydrogen, p is 2, R^1 is hydrogen and R^2 is tert-butyl and the combination of B, D, E and R^b for a compound corresponds in each case to a row of Table A.

Table 10

- 25 Compounds of the formula I.5 wherein X is hydrogen, p is 2, R^1 is hydrogen and R^2 is phenyl and the combination of B, D, E and R^c for a compound corresponds in each case to a row of Table A.

Table 11

- 30 Compounds of the formula I.5 wherein X is hydrogen, p is 4, R^1 and R^2 are hydrogen and the combination of B, D, E and R^c for a compound corresponds in each case to a row of Table A.

Table 12

- 35 Compounds of the formula I.5 wherein X is hydrogen, p is 4, R^1 is hydrogen and R^2 is methyl and the combination of B, D, E and R^c for a compound corresponds in each case to a row of Table A.

Table 13

- 40 Compounds of the formula I.5 wherein X is hydrogen, p is 4, R^1 is hydrogen and R^2 is ethyl and the combination of B, D, E and R^c for a compound corresponds in each case to a row of Table A.

Table 14

- 45 Compounds of the formula I.5 wherein X is hydrogen, p is 4, R^1 is hydrogen and R^2 is i-propyl and the combination of B, D, E and R^c

17

for a compound corresponds in each case to a row of Table A.

Table 15

Compounds of the formula I.5 wherein X is hydrogen, p is 4, R¹ is
5 hydrogen and R² is n-propyl and the combination of B, D, E and R^c
for a compound corresponds in each case to a row of Table A.

Table 16

Compounds of the formula I.5 wherein X is hydrogen, p is 4, R¹ is
10 hydrogen and R² is cyclo-propyl and the combination of B, D, E and
R^c for a compound corresponds in each case to a row of Table A.

Table 17

Compounds of the formula I.5 wherein X is hydrogen, p is 4, R¹ is
15 hydrogen and R² is n-butyl and the combination of B, D, E and R^c
for a compound corresponds in each case to a row of Table A.

Table 18

Compounds of the formula I.5 wherein X is hydrogen, p is 4, R¹ is
20 hydrogen and R² is i-butyl and the combination of B, D, E and R^c
for a compound corresponds in each case to a row of Table A.

Table 19

Compounds of the formula I.5 wherein X is hydrogen, p is 4, R¹ is
25 hydrogen and R² is tert-butyl and the combination of B, D, E and
R^c for a compound corresponds in each case to a row of Table A.

Table 20

Compounds of the formula I.5 wherein X is hydrogen, p is 4, R¹ is
30 hydrogen and R² is phenyl and the combination of B, D, E and R^c
for a compound corresponds in each case to a row of Table A.

Table 21

Compounds of the formula I.5 wherein X is fluorine, p is 2, R¹ and
35 R² are hydrogen and the combination of B, D, E and R^c for a
compound corresponds in each case to a row of Table A.

Table 22

Compounds of the formula I.5 wherein X is fluorine, p is 2, R¹ is
40 hydrogen and R² is methyl and the combination of B, D, E and R^c
for a compound corresponds in each case to a row of Table A.

Table 23

Compounds of the formula I.5 wherein X is fluorine, p is 2, R¹ is
45 hydrogen and R² is ethyl and the combination of B, D, E and R^c for
a compound corresponds in each case to a row of Table A.

Table 24

Compounds of the formula I.5 wherein X is fluorine, p is 2, R¹ is hydrogen and R² is i-propyl and the combination of B, D, E and R^c for a compound corresponds in each case to a row of Table A.

5

Table 25

Compounds of the formula I.5 wherein X is fluorine, p is 2, R¹ is hydrogen and R² is n-propyl and the combination of B, D, E and R^c for a compound corresponds in each case to a row of Table A.

10

Table 26

Compounds of the formula I.5 wherein X is fluorine, p is 2, R¹ is hydrogen and R² is cyclo-propyl and the combination of B, D, E and R^c for a compound corresponds in each case to a row of Table A.

15

Table 27

Compounds of the formula I.5 wherein X is fluorine, p is 2, R¹ is hydrogen and R² is n-butyl and the combination of B, D, E and R^c for a compound corresponds in each case to a row of Table A.

20

Table 28

Compounds of the formula I.5 wherein X is fluorine, p is 2, R¹ is hydrogen and R² is i-butyl and the combination of B, D, E and R^c for a compound corresponds in each case to a row of Table A.

25

Table 29

Compounds of the formula I.5 wherein X is fluorine, p is 2, R¹ is hydrogen and R² is tert-butyl and the combination of B, D, E and R^c for a compound corresponds in each case to a row of Table A.

30

Table 30

Compounds of the formula I.5 wherein X is fluorine, p is 2, R¹ is hydrogen and R² is phenyl and the combination of B, D, E and R^c for a compound corresponds in each case to a row of Table A.

35

Table 31

Compounds of the formula I.5 wherein X is fluorine, p is 4, R¹ and R² are hydrogen and the combination of B, D, E and R^c for a compound corresponds in each case to a row of Table A.

40

Table 32

Compounds of the formula I.5 wherein X is fluorine, p is 4, R¹ is hydrogen and R² is methyl and the combination of B, D, E and R^c for a compound corresponds in each case to a row of Table A.

45

Table 33

Compounds of the formula I.5 wherein X is fluorine, p is 4, R¹ is

19

hydrogen and R² is ethyl and the combination of B, D, E and R^c for a compound corresponds in each case to a row of Table A.

Table 34

- 5 Compounds of the formula I.5 wherein X is fluorine, p is 4, R¹ is hydrogen and R² is i-propyl and the combination of B, D, E and R^c for a compound corresponds in each case to a row of Table A.

Table 35

- 10 Compounds of the formula I.5 wherein X is fluorine, p is 4, R¹ is hydrogen and R² is n-propyl and the combination of B, D, E and R^c for a compound corresponds in each case to a row of Table A.

Table 36

- 15 Compounds of the formula I.5 wherein X is fluorine, p is 4, R¹ is hydrogen and R² is cyclo-propyl and the combination of B, D, E and R^c for a compound corresponds in each case to a row of Table A.

Table 37

- 20 Compounds of the formula I.5 wherein X is fluorine, p is 4, R¹ is hydrogen and R² is n-butyl and the combination of B, D, E and R^c for a compound corresponds in each case to a row of Table A.

Table 38

- 25 Compounds of the formula I.5 wherein X is fluorine, p is 4, R¹ is hydrogen and R² is i-butyl and the combination of B, D, E and R^c for a compound corresponds in each case to a row of Table A.

Table 39


- 30 Compounds of the formula I.5 wherein X is fluorine, p is 4, R¹ is hydrogen and R² is tert-butyl and the combination of B, D, E and R^c for a compound corresponds in each case to a row of Table A.

Table 40

- 35 Compounds of the formula I.5 wherein X is fluorine, p is 4, R¹ is hydrogen and R² is phenyl and the combination of B, D, E and R^c for a compound corresponds in each case to a row of Table A.

Table A

40



45

5	Nr.	B	D	E	R^c
	A-1	CH	CH	O	-
	A-2	CH	CH	O	4-CH ₃
	A-3	CH	CH	O	4-CH ₂ CH ₃
10	A-4	CH	CH	O	4-CH ₂ CH ₂ CH ₃
	A-5	CH	CH	O	4-CH(CH ₃) ₂
	A-6	CH	CH	O	4-CH ₂ CH ₂ CH ₂ CH ₃
	A-7	CH	CH	O	4-CH ₂ CH(CH ₃) ₂
	A-8	CH	CH	O	4-CH(CH ₃) ₃
15	A-9	CH	CH	O	4-F
	A-10	CH	CH	O	4-Cl
	A-11	CH	CH	O	4-Br
	A-12	CH	CH	O	4-I
20	A-13	CH	CH	O	4-CF ₃
	A-14	CH	CH	O	4-OCF ₃
	A-15	CH	CH	O	4-OCH ₃
	A-16	CH	CH	O	4-OCH ₂ CH ₃
25	A-17	CH	CH	O	4-OCH ₂ CH ₂ CH ₃
	A-18	CH	CH	O	4-OCH(CH ₃) ₂
	A-19	CH	CH	O	4-NO ₂
	A-20	CH	CH	O	4-NH ₂
	A-21	CH	CH	O	4-NH(C=O)CH ₃
30	A-22	CH	CH	O	5-CH ₃
	A-23	CH	CH	O	5-CH ₂ CH ₃
	A-24	CH	CH	O	5-CH ₂ CH ₂ CH ₃
	A-25	CH	CH	O	5-CH(CH ₃) ₂
35	A-26	CH	CH	O	5-CH ₂ CH ₂ CH ₂ CH ₃
	A-27	CH	CH	O	5-CH ₂ CH(CH ₃) ₂
	A-28	CH	CH	O	5-CH(CH ₃) ₃
	A-29	CH	CH	O	5-F
40	A-30	CH	CH	O	5-Cl
	A-31	CH	CH	O	5-Br
	A-32	CH	CH	O	5-I
	A-33	CH	CH	O	5-CF ₃
45	A-34	CH	CH	O	5-OCF ₃
	A-35	CH	CH	O	5-OCH ₃
	A-36	CH	CH	O	5-OCH ₂ CH ₃

21

	Nr.	B	D	E	R ^a
5	A-37	CH	CH	O	5-OCH ₂ CH ₂ CH ₃
	A-38	CH	CH	O	5-OCH(CH ₃) ₂
	A-39	CH	CH	O	5-NO ₂
	A-40	CH	CH	O	5-NH ₂
	A-41	CH	CH	O	5-NH(C=O)CH ₃
10	A-42	CH	CH	O	6-CH ₃
	A-43	CH	CH	O	6-CH ₂ CH ₃
	A-44	CH	CH	O	6-CH ₂ CH ₂ CH ₃
	A-45	CH	CH	O	6-CH(CH ₃) ₂
	A-46	CH	CH	O	6-CH ₂ CH ₂ CH ₂ CH ₃
15	A-47	CH	CH	O	6-CH ₂ CH(CH ₃) ₂
	A-48	CH	CH	O	6-CH(CH ₃) ₃
	A-49	CH	CH	O	6-F
	A-50	CH	CH	O	6-Cl
	A-51	CH	CH	O	6-Br
20	A-52	CH	CH	O	6-I
	A-53	CH	CH	O	6-CF ₃
	A-54	CH	CH	O	6-OCF ₃
	A-55	CH	CH	O	6-OCH ₃
	A-56	CH	CH	O	6-OCH ₂ CH ₃
25	A-57	CH	CH	O	6-OCH ₂ CH ₂ CH ₃
	A-58	CH	CH	O	6-OCH(CH ₃) ₂
	A-59	CH	CH	O	6-NO ₂
	A-60	CH	CH	O	6-NH ₂
	A-61	CH	CH	O	6-NH(C=O)CH ₃
30	A-62	CH	CH	O	7-CH ₃
	A-63	CH	CH	O	7-CH ₂ CH ₃
	A-64	CH	CH	O	7-CH ₂ CH ₂ CH ₃
	A-65	CH	CH	O	7-CH(CH ₃) ₂
	A-66	CH	CH	O	7-CH ₂ CH ₂ CH ₂ CH ₃
35	A-67	CH	CH	O	7-CH ₂ CH(CH ₃) ₂
	A-68	CH	CH	O	7-CH(CH ₃) ₃
	A-69	CH	CH	O	7-F
	A-70	CH	CH	O	7-Cl
	A-71	CH	CH	O	7-Br
40	A-72	CH	CH	O	7-I
	A-73	CH	CH	O	7-CF ₃
	A-74	CH	CH	O	7-OCF ₃
	A-75	CH	CH	O	7-OCH ₃
	A-75	CH	CH	O	7-OCH ₃

22

	Nr.	B	D	E	R ^c
5	A-76	CH	CH	O	7-OCH ₂ CH ₃
	A-77	CH	CH	O	7-OCH ₂ CH ₂ CH ₃
	A-78	CH	CH	O	7-OCH(CH ₃) ₂
	A-79	CH	CH	O	7-NO ₂
	A-80	CH	CH	O	7-NH ₂
10	A-81	CH	CH	O	7-NH(C=O)CH ₃
	A-82	CH	CH	S	-
	A-83	CH	CH	S	4-CH ₃
	A-84	CH	CH	S	4-CH ₂ CH ₃
	A-85	CH	CH	S	4-CH ₂ CH ₂ CH ₃
15	A-86	CH	CH	S	4-CH(CH ₃) ₂
	A-87	CH	CH	S	4-CH ₂ CH ₂ CH ₂ CH ₃
	A-88	CH	CH	S	4-CH ₂ CH(CH ₃) ₂
	A-89	CH	CH	S	4-CH(CH ₃) ₃
	A-90	CH	CH	S	4-F
20	A-91	CH	CH	S	4-Cl
	A-92	CH	CH	S	4-Br
	A-93	CH	CH	S	4-I
	A-94	CH	CH	S	4-CF ₃
	A-95	CH	CH	S	4-OCF ₃
25	A-96	CH	CH	S	4-OCH ₃
	A-97	CH	CH	S	4-OCH ₂ CH ₃
	A-98	CH	CH	S	4-OCH ₂ CH ₂ CH ₃
	A-99	CH	CH	S	4-OCH(CH ₃) ₂
	A-100	CH	CH	S	4-NO ₂
30	A-101	CH	CH	S	4-NH ₂
	A-102	CH	CH	S	4-NH(C=O)CH ₃
	A-103	CH	CH	S	5-CH ₃
	A-104	CH	CH	S	5-CH ₂ CH ₃
	A-105	CH	CH	S	5-CH ₂ CH ₂ CH ₃
35	A-106	CH	CH	S	5-CH(CH ₃) ₂
	A-107	CH	CH	S	5-CH ₂ CH ₂ CH ₂ CH ₃
	A-108	CH	CH	S	5-CH ₂ CH(CH ₃) ₂
	A-109	CH	CH	S	5-CH(CH ₃) ₃
	A-110	CH	CH	S	5-F
40	A-111	CH	CH	S	5-Cl
	A-112	CH	CH	S	5-Br
	A-113	CH	CH	S	5-I
	A-114	CH	CH	S	5-CF ₃
	A-114	CH	CH	S	5-CF ₃

23

	Nr.	B	D	E	R ^c
5	A-115	CH	CH	S	5-OCF ₃
	A-116	CH	CH	S	5-OCH ₃
	A-117	CH	CH	S	5-OCH ₂ CH ₃
	A-118	CH	CH	S	5-OCH ₂ CH ₂ CH ₃
	A-119	CH	CH	S	5-OCH(CH ₃) ₂
10	A-120	CH	CH	S	5-NO ₂
	A-121	CH	CH	S	5-NH ₂
	A-122	CH	CH	S	5-NH(C=O)CH ₃
	A-123	CH	CH	S	6-CH ₃
	A-124	CH	CH	S	6-CH ₂ CH ₃
15	A-125	CH	CH	S	6-CH ₂ CH ₂ CH ₃
	A-126	CH	CH	S	6-CH(CH ₃) ₂
	A-127	CH	CH	S	6-CH ₂ CH ₂ CH ₂ CH ₃
	A-128	CH	CH	S	6-CH ₂ CH(CH ₃) ₂
	A-129	CH	CH	S	6-CH(CH ₃) ₃
20	A-130	CH	CH	S	6-F
	A-131	CH	CH	S	6-Cl
	A-132	CH	CH	S	6-Br
	A-133	CH	CH	S	6-I
	A-134	CH	CH	S	6-CF ₃
25	A-135	CH	CH	S	6-OCF ₃
	A-136	CH	CH	S	6-OCH ₃
	A-137	CH	CH	S	6-OCH ₂ CH ₃
	A-138	CH	CH	S	6-OCH ₂ CH ₂ CH ₃
	A-139	CH	CH	S	6-OCH(CH ₃) ₂
30	A-140	CH	CH	S	6-NO ₂
	A-141	CH	CH	S	6-NH ₂
	A-142	CH	CH	S	6-NH(C=O)CH ₃
	A-143	CH	CH	S	7-CH ₃
	A-144	CH	CH	S	7-CH ₂ CH ₃
35	A-145	CH	CH	S	7-CH ₂ CH ₂ CH ₃
	A-146	CH	CH	S	7-CH(CH ₃) ₂
	A-147	CH	CH	S	7-CH ₂ CH ₂ CH ₂ CH ₃
	A-148	CH	CH	S	7-CH ₂ CH(CH ₃) ₂
	A-149	CH	CH	S	7-CH(CH ₃) ₃
40	A-150	CH	CH	S	7-F
	A-151	CH	CH	S	7-Cl
	A-152	CH	CH	S	7-Br
	A-153	CH	CH	S	7-I

	Nr.	B	D	E	R ^c
5	A-154	CH	CH	S	7-CF ₃
	A-155	CH	CH	S	7-OCF ₃
	A-156	CH	CH	S	7-OCH ₃
	A-157	CH	CH	S	7-OCH ₂ CH ₃
	A-158	CH	CH	S	7-OCH ₂ CH ₂ CH ₃
10	A-159	CH	CH	S	7-OCH(CH ₃) ₂
	A-160	CH	CH	S	7-NO ₂
	A-161	CH	CH	S	7-NH ₂
	A-162	CH	CH	S	7-NH(C=O)CH ₃
	A-163	CH	N	O	-
15	A-164	CH	N	O	4-CH ₃
	A-165	CH	N	O	4-CH ₂ CH ₃
	A-166	CH	N	O	4-CH ₂ CH ₂ CH ₃
	A-167	CH	N	O	4-CH(CH ₃) ₂
	A-168	CH	N	O	4-CH ₂ CH ₂ CH ₂ CH ₃
20	A-169	CH	N	O	4-CH ₂ CH(CH ₃) ₂
	A-170	CH	N	O	4-CH(CH ₃) ₃
	A-171	CH	N	O	4-F
	A-172	CH	N	O	4-Cl
	A-173	CH	N	O	4-Br
25	A-174	CH	N	O	4-I
	A-175	CH	N	O	4-CF ₃
	A-176	CH	N	O	4-OCF ₃
	A-177	CH	N	O	4-OCH ₃
	A-178	CH	N	O	4-OCH ₂ CH ₃
30	A-179	CH	N	O	4-OCH ₂ CH ₂ CH ₃
	A-180	CH	N	O	4-OCH(CH ₃) ₂
	A-181	CH	N	O	4-NO ₂
	A-182	CH	N	O	4-NH ₂
	A-183	CH	N	O	4-NH(C=O)CH ₃
35	A-184	CH	N	O	5-CH ₃
	A-185	CH	N	O	5-CH ₂ CH ₃
	A-186	CH	N	O	5-CH ₂ CH ₂ CH ₃
	A-187	CH	N	O	5-CH(CH ₃) ₂
	A-188	CH	N	O	5-CH ₂ CH ₂ CH ₂ CH ₃
40	A-189	CH	N	O	5-CH ₂ CH(CH ₃) ₂
	A-190	CH	N	O	5-CH(CH ₃) ₃
	A-191	CH	N	O	5-F
	A-192	CH	N	O	5-Cl

25

	Nr.	B	D	E	R ^c
5	A-193	CH	N	O	5-Br
	A-194	CH	N	O	5-I
	A-195	CH	N	O	5-CF ₃
	A-196	CH	N	O	5-OCF ₃
	A-197	CH	N	O	5-OCH ₃
10	A-198	CH	N	O	5-OCH ₂ CH ₃
	A-199	CH	N	O	5-OCH ₂ CH ₂ CH ₃
	A-200	CH	N	O	5-OCH(CH ₃) ₂
	A-201	CH	N	O	5-NO ₂
	A-202	CH	N	O	5-NH ₂
15	A-203	CH	N	O	5-NH(C=O)CH ₃
	A-204	CH	N	O	6-CH ₃
	A-205	CH	N	O	6-CH ₂ CH ₃
	A-206	CH	N	O	6-CH ₂ CH ₂ CH ₃
	A-207	CH	N	O	6-CH(CH ₃) ₂
20	A-208	CH	N	O	6-CH ₂ CH ₂ CH ₂ CH ₃
	A-209	CH	N	O	6-CH ₂ CH(CH ₃) ₂
	A-210	CH	N	O	6-CH(CH ₃) ₃
	A-211	CH	N	O	6-F
	A-212	CH	N	O	6-Cl
25	A-213	CH	N	O	6-Br
	A-214	CH	N	O	6-I
	A-215	CH	N	O	6-CF ₃
	A-216	CH	N	O	6-OCF ₃
	A-217	CH	N	O	6-OCH ₃
30	A-218	CH	N	O	6-OCH ₂ CH ₃
	A-219	CH	N	O	6-OCH ₂ CH ₂ CH ₃
	A-220	CH	N	O	6-OCH(CH ₃) ₂
	A-221	CH	N	O	6-NO ₂
	A-222	CH	N	O	6-NH ₂
35	A-223	CH	N	O	6-NH(C=O)CH ₃
	A-224	CH	N	S	-
	A-225	CH	N	S	4-CH ₃
	A-226	CH	N	S	4-CH ₂ CH ₃
	A-227	CH	N	S	4-CH ₂ CH ₂ CH ₃
40	A-228	CH	N	S	4-CH(CH ₃) ₂
	A-229	CH	N	S	4-CH ₂ CH ₂ CH ₂ CH ₃
	A-230	CH	N	S	4-CH ₂ CH(CH ₃) ₂
	A-231	CH	N	S	4-CH(CH ₃) ₃
	A-231	CH	N	S	4-CH(CH ₃) ₃

26

	Nr.	B	D	E	R ^c
5	A-232	CH	N	S	4-F
	A-233	CH	N	S	4-Cl
	A-234	CH	N	S	4-Br
	A-235	CH	N	S	4-I
	A-236	CH	N	S	4-CF ₃
10	A-237	CH	N	S	4-OCF ₃
	A-238	CH	N	S	4-OCH ₃
	A-239	CH	N	S	4-OCH ₂ CH ₃
	A-240	CH	N	S	4-OCH ₂ CH ₂ CH ₃
	A-241	CH	N	S	4-OCH(CH ₃) ₂
15	A-242	CH	N	S	4-NO ₂
	A-243	CH	N	S	4-NH ₂
	A-244	CH	N	S	4-NH(C=O)CH ₃
	A-245	CH	N	S	5-CH ₃
	A-246	CH	N	S	5-CH ₂ CH ₃
20	A-247	CH	N	S	5-CH ₂ CH ₂ CH ₃
	A-248	CH	N	S	5-CH(CH ₃) ₂
	A-249	CH	N	S	5-CH ₂ CH ₂ CH ₂ CH ₃
	A-250	CH	N	S	5-CH ₂ CH(CH ₃) ₂
	A-251	CH	N	S	5-CH(CH ₃) ₃
25	A-252	CH	N	S	5-F
	A-253	CH	N	S	5-Cl
	A-254	CH	N	S	5-Br
	A-255	CH	N	S	5-I
	A-256	CH	N	S	5-CF ₃
30	A-257	CH	N	S	5-OCF ₃
	A-258	CH	N	S	5-OCH ₃
	A-259	CH	N	S	5-OCH ₂ CH ₃
	A-260	CH	N	S	5-OCH ₂ CH ₂ CH ₃
	A-261	CH	N	S	5-OCH(CH ₃) ₂
35	A-262	CH	N	S	5-NO ₂
	A-263	CH	N	S	5-NH ₂
	A-264	CH	N	S	5-NH(C=O)CH ₃
	A-265	CH	N	S	6-CH ₃
	A-266	CH	N	S	6-CH ₂ CH ₃
40	A-267	CH	N	S	6-CH ₂ CH ₂ CH ₃
	A-268	CH	N	S	6-CH(CH ₃) ₂
	A-269	CH	N	S	6-CH ₂ CH ₂ CH ₂ CH ₃
	A-270	CH	N	S	6-CH ₂ CH(CH ₃) ₂

	Nr.	B	D	E	R ^c
5	A-271	CH	N	S	6-CH(CH ₃) ₃
	A-272	CH	N	S	6-F
	A-273	CH	N	S	6-Cl
	A-274	CH	N	S	6-Br
	A-275	CH	N	S	6-I
10	A-276	CH	N	S	6-CF ₃
	A-277	CH	N	S	6-OCF ₃
	A-278	CH	N	S	6-OCH ₃
	A-279	CH	N	S	6-OCH ₂ CH ₃
	A-280	CH	N	S	6-OCH ₂ CH ₂ CH ₃
15	A-281	CH	N	S	6-OCH(CH ₃) ₂
	A-282	CH	N	S	6-NO ₂
	A-283	CH	N	S	6-NH ₂
	A-284	CH	N	S	6-NH(C=O)CH ₃
	A-285	N	CH	O	-
20	A-286	N	CH	O	5-CH ₃
	A-287	N	CH	O	5-CH ₂ CH ₃
	A-288	N	CH	O	5-CH ₂ CH ₂ CH ₃
	A-289	N	CH	O	5-CH(CH ₃) ₂
	A-290	N	CH	O	5-CH ₂ CH ₂ CH ₂ CH ₃
25	A-291	N	CH	O	5-CH ₂ CH(CH ₃) ₂
	A-292	N	CH	O	5-CH(CH ₃) ₃
	A-293	N	CH	O	5-F
	A-294	N	CH	O	5-Cl
	A-295	N	CH	O	5-Br
30	A-296	N	CH	O	5-I
	A-297	N	CH	O	5-CF ₃
	A-298	N	CH	O	5-OCF ₃
	A-299	N	CH	O	5-OCH ₃
	A-300	N	CH	O	5-OCH ₂ CH ₃
35	A-301	N	CH	O	5-OCH ₂ CH ₂ CH ₃
	A-302	N	CH	O	5-OCH(CH ₃) ₂
	A-303	N	CH	O	5-NO ₂
	A-304	N	CH	O	5-NH ₂
	A-305	N	CH	O	5-NH(C=O)CH ₃
40	A-306	N	CH	O	6-CH ₃
	A-307	N	CH	O	6-CH ₂ CH ₃
	A-308	N	CH	O	6-CH ₂ CH ₂ CH ₃
	A-309	N	CH	O	6-CH(CH ₃) ₂


28

	Nr.	B	D	E	R ^c
5	A-310	N	CH	O	6-CH ₂ CH ₂ CH ₂ CH ₃
	A-311	N	CH	O	6-CH ₂ CH(CH ₃) ₂
	A-312	N	CH	O	6-CH(CH ₃) ₃
	A-313	N	CH	O	6-F
	A-314	N	CH	O	6-Cl
10	A-315	N	CH	O	6-Br
	A-316	N	CH	O	6-I
	A-317	N	CH	O	6-CF ₃
	A-318	N	CH	O	6-OCF ₃
	A-319	N	CH	O	6-OCH ₃
15	A-320	N	CH	O	6-OCH ₂ CH ₃
	A-321	N	CH	O	6-OCH ₂ CH ₂ CH ₃
	A-322	N	CH	O	6-OCH(CH ₃) ₂
	A-323	N	CH	O	6-NO ₂
	A-324	N	CH	O	6-NH ₂
20	A-325	N	CH	O	6-NH(C=O)CH ₃
	A-326	N	CH	O	7-CH ₃
	A-327	N	CH	O	7-CH ₂ CH ₃
	A-328	N	CH	O	7-CH ₂ CH ₂ CH ₃
	A-329	N	CH	O	7-CH(CH ₃) ₂
25	A-330	N	CH	O	7-CH ₂ CH ₂ CH ₂ CH ₃
	A-331	N	CH	O	7-CH ₂ CH(CH ₃) ₂
	A-332	N	CH	O	7-CH(CH ₃) ₃
	A-333	N	CH	O	7-F
	A-334	N	CH	O	7-Cl
30	A-335	N	CH	O	7-Br
	A-336	N	CH	O	7-I
	A-337	N	CH	O	7-CF ₃
	A-338	N	CH	O	7-OCF ₃
	A-339	N	CH	O	7-OCH ₃
35	A-340	N	CH	O	7-OCH ₂ CH ₃
	A-341	N	CH	O	7-OCH ₂ CH ₂ CH ₃
	A-342	N	CH	O	7-OCH(CH ₃) ₂
	A-343	N	CH	O	7-NO ₂
	A-344	N	CH	O	7-NH ₂
40	A-345	N	CH	O	7-NH(C=O)CH ₃

45

Table 41

Compounds of the formula I.6



wherein X is hydrogen, p is 2, R¹ and R² are hydrogen, E' is oxygen and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

10

Table 42

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is methyl, E' is oxygen and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

15

Table 43

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is ethyl, E' is oxygen and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

20

Table 44

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is i-propyl, E' is oxygen and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

25

Table 45

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is n-propyl, E' is oxygen and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

30

Table 46

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is cyclo-propyl, E' is oxygen and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

35

Table 47

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is n-butyl, E' is oxygen and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

40

Table 48

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is i-butyl, E' is oxygen and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

45

Table 49

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is

30

hydrogen and R² is tert-butyl, E' is oxygen and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

Table 50

- 5 Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is phenyl, E' is oxygen and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

Table 51

- 10 Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ and R² are hydrogen, E' is oxygen and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

Table 52

- 15 Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is hydrogen and R² is methyl, E' is oxygen and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

Table 53

- 20 Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is hydrogen and R² is ethyl, E' is oxygen and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

Table 54

- 25 Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is hydrogen and R² is i-propyl, E' is oxygen and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

Table 55

- 30 Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is hydrogen and R² is n-propyl, E' is oxygen and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

Table 56

- 35 Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is hydrogen and R² is cyclo-propyl, E' is oxygen and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

Table 57

- 40 Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is hydrogen and R² is n-butyl, E' is oxygen and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

Table 58

- 45 Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is hydrogen and R² is i-butyl, E' is oxygen and (R^c)_q is in the

31

4-position and corresponds in each case to a row of Table B.

Table 59

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is
5 hydrogen and R² is tert-butyl, E' is oxygen and (R^c)_q is in the
4-position and corresponds in each case to a row of Table B.

Table 60

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is
10 hydrogen and R² is phenyl, E' is oxygen and (R^c)_q is in the
4-position and corresponds in each case to a row of Table B.

Table 61

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ and
15 R² are hydrogen, E' is oxygen and (R^c)_q is in the 4-position and
corresponds in each case to a row of Table B.

Table 62

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is
20 hydrogen and R² is methyl, E' is oxygen and (R^c)_q is in the
4-position and corresponds in each case to a row of Table B.

Table 63

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is
25 hydrogen and R² is ethyl, E' is oxygen and (R^c)_q is in the
4-position and corresponds in each case to a row of Table B.

Table 64

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is
30 hydrogen and R² is i-propyl, E' is oxygen and (R^c)_q is in the
4-position and corresponds in each case to a row of Table B.

Table 65

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is
35 hydrogen and R² is n-propyl, E' is oxygen and (R^c)_q is in the
4-position and corresponds in each case to a row of Table B.

Table 66

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is
40 hydrogen and R² is cyclo-propyl, E' is oxygen and (R^c)_q is in the
4-position and corresponds in each case to a row of Table B.

Table 67

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is
45 hydrogen and R² is n-butyl, E' is oxygen and (R^c)_q is in the
4-position and corresponds in each case to a row of Table B.

Table 68

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is hydrogen and R² is i-butyl, E' is oxygen and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

5

Table 69

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is hydrogen and R² is tert-butyl, E' is oxygen and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

10

Table 70

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is hydrogen and R² is phenyl, E' is oxygen and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

15

Table 71

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ and R² are hydrogen, E' is oxygen and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

20

Table 72

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ is hydrogen and R² is methyl, E' is oxygen and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

25

Table 73

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ is hydrogen and R² is ethyl, E' is oxygen and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

30

Table 74

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ is hydrogen and R² is i-propyl, E' is oxygen and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

35

Table 75

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ is hydrogen and R² is n-propyl, E' is oxygen and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

40

Table 76

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ is hydrogen and R² is cyclo-propyl, E' is oxygen and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

45

Table 77

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ is

33

hydrogen and R^2 is n-butyl, E' is oxygen and $(R^c)_q$ is in the 4-position and corresponds in each case to a row of Table B.

Table 78

- 5 Compounds of the formula I.6 wherein X is fluorine, p is 4, R^1 is hydrogen and R^2 is i-butyl, E' is oxygen and $(R^c)_q$ is in the 4-position and corresponds in each case to a row of Table B.

Table 79

- 10 Compounds of the formula I.6 wherein X is fluorine, p is 4, R^1 is hydrogen and R^2 is tert-butyl, E' is oxygen and $(R^c)_q$ is in the 4-position and corresponds in each case to a row of Table B.

Table 80

- 15 Compounds of the formula I.6 wherein X is fluorine, p is 4, R^1 is hydrogen and R^2 is phenyl, E' is oxygen and $(R^c)_q$ is in the 4-position and corresponds in each case to a row of Table B.

Table 81

- 20 Compounds of the formula I.6 wherein X is hydrogen, p is 2, R^1 and R^2 are hydrogen, E' is oxygen and $(R^c)_q$ is in the 5-position and corresponds in each case to a row of Table B.

Table 82

- 25 Compounds of the formula I.6 wherein X is hydrogen, p is 2, R^1 is hydrogen and R^2 is methyl, E' is oxygen and $(R^c)_q$ is in the 5-position and corresponds in each case to a row of Table B.

Table 83

- 30 Compounds of the formula I.6 wherein X is hydrogen, p is 2, R^1 is hydrogen and R^2 is ethyl, E' is oxygen and $(R^c)_q$ is in the 5-position and corresponds in each case to a row of Table B.

Table 84

- 35 Compounds of the formula I.6 wherein X is hydrogen, p is 2, R^1 is hydrogen and R^2 is i-propyl, E' is oxygen and $(R^c)_q$ is in the 5-position and corresponds in each case to a row of Table B.

Table 85

- 40 Compounds of the formula I.6 wherein X is hydrogen, p is 2, R^1 is hydrogen and R^2 is n-propyl, E' is oxygen and $(R^c)_q$ is in the 5-position and corresponds in each case to a row of Table B.

Table 86

- 45 Compounds of the formula I.6 wherein X is hydrogen, p is 2, R^1 is hydrogen and R^2 is cyclo-propyl, E' is oxygen and $(R^c)_q$ is in the

34

5-position and corresponds in each case to a row of Table B.

Table 87

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is
5 hydrogen and R² is n-butyl, E' is oxygen and (R^c)_q is in the
5-position and corresponds in each case to a row of Table B.

Table 88

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is
10 hydrogen and R² is i-butyl, E' is oxygen and (R^c)_q is in the
5-position and corresponds in each case to a row of Table B.

Table 89

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is
15 hydrogen and R² is tert-butyl, E' is oxygen and (R^c)_q is in the
5-position and corresponds in each case to a row of Table B.

Table 90

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is
20 hydrogen and R² is phenyl, E' is oxygen and (R^c)_q is in the
5-position and corresponds in each case to a row of Table B.

Table 91

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ and
25 R² are hydrogen, E' is oxygen and (R^c)_q is in the 5-position and
corresponds in each case to a row of Table B.

Table 92

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is
30 hydrogen and R² is methyl, E' is oxygen and (R^c)_q is in the
5-position and corresponds in each case to a row of Table B.

Table 93

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is
35 hydrogen and R² is ethyl, E' is oxygen and (R^c)_q is in the
5-position and corresponds in each case to a row of Table B.

Table 94

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is
40 hydrogen and R² is i-propyl, E' is oxygen and (R^c)_q is in the
5-position and corresponds in each case to a row of Table B.

Table 95

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is
45 hydrogen and R² is n-propyl, E' is oxygen and (R^c)_q is in the
5-position and corresponds in each case to a row of Table B.

35

Table 96

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is hydrogen and R² is cyclo-propyl, E' is oxygen and (R^c)_q is in the 5-position and corresponds in each case to a row of Table B.

5

Table 97

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is hydrogen and R² is n-butyl, E' is oxygen and (R^c)_q is in the 5-position and corresponds in each case to a row of Table B.

10

Table 98

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is hydrogen and R² is i-butyl, E' is oxygen and (R^c)_q is in the 5-position and corresponds in each case to a row of Table B.

15

Table 99

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is hydrogen and R² is tert-butyl, E' is oxygen and (R^c)_q is in the 5-position and corresponds in each case to a row of Table B.

20

Table 100

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is hydrogen and R² is phenyl, E' is oxygen and (R^c)_q is in the 5-position and corresponds in each case to a row of Table B.

25

Table 101

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ and R² are hydrogen, E' is oxygen and (R^c)_q is in the 5-position and corresponds in each case to a row of Table B.

30

Table 102

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is hydrogen and R² is methyl, E' is oxygen and (R^c)_q is in the 5-position and corresponds in each case to a row of Table B.

35

Table 103

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is hydrogen and R² is ethyl, E' is oxygen and (R^c)_q is in the 5-position and corresponds in each case to a row of Table B.

40

Table 104

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is hydrogen and R² is i-propyl, E' is oxygen and (R^c)_q is in the 5-position and corresponds in each case to a row of Table B.

45

Table 105

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is

36

hydrogen and R^2 is n-propyl, E' is oxygen and $(R^c)_q$ is in the 5-position and corresponds in each case to a row of Table B.

Table 106

- 5 Compounds of the formula I.6 wherein X is fluorine, p is 2, R^1 is hydrogen and R^2 is cyclo-propyl, E' is oxygen and $(R^c)_q$ is in the 5-position and corresponds in each case to a row of Table B.

Table 107

- 10 Compounds of the formula I.6 wherein X is fluorine, p is 2, R^1 is hydrogen and R^2 is n-butyl, E' is oxygen and $(R^c)_q$ is in the 5-position and corresponds in each case to a row of Table B.

Table 108

- 15 Compounds of the formula I.6 wherein X is fluorine, p is 2, R^1 is hydrogen and R^2 is i-butyl, E' is oxygen and $(R^c)_q$ is in the 5-position and corresponds in each case to a row of Table B.

Table 109

- 20 Compounds of the formula I.6 wherein X is fluorine, p is 2, R^1 is hydrogen and R^2 is tert-butyl, E' is oxygen and $(R^c)_q$ is in the 5-position and corresponds in each case to a row of Table B.

Table 110

- 25 Compounds of the formula I.6 wherein X is fluorine, p is 2, R^1 is hydrogen and R^2 is phenyl, E' is oxygen and $(R^c)_q$ is in the 5-position and corresponds in each case to a row of Table B.

Table 111

- 30 Compounds of the formula I.6 wherein X is fluorine, p is 4, R^1 and R^2 are hydrogen, E' is oxygen and $(R^c)_q$ is in the 5-position and corresponds in each case to a row of Table B.

Table 112

- 35 Compounds of the formula I.6 wherein X is fluorine, p is 4, R^1 is hydrogen and R^2 is methyl, E' is oxygen and $(R^c)_q$ is in the 5-position and corresponds in each case to a row of Table B.

Table 113

- 40 Compounds of the formula I.6 wherein X is fluorine, p is 4, R^1 is hydrogen and R^2 is ethyl, E' is oxygen and $(R^c)_q$ is in the 5-position and corresponds in each case to a row of Table B.

Table 114

- 45 Compounds of the formula I.6 wherein X is fluorine, p is 4, R^1 is hydrogen and R^2 is i-propyl, E' is oxygen and $(R^c)_q$ is in the

37

5-position and corresponds in each case to a row of Table B.

Table 115

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ is 5 hydrogen and R² is n-propyl, E' is oxygen and (R^c)_q is in the 5-position and corresponds in each case to a row of Table B.

Table 116

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ is 10 hydrogen and R² is cyclo-propyl, E' is oxygen and (R^c)_q is in the 5-position and corresponds in each case to a row of Table B.

Table 117

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ is 15 hydrogen and R² is n-butyl, E' is oxygen and (R^c)_q is in the 5-position and corresponds in each case to a row of Table B.

Table 118

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ is 20 hydrogen and R² is i-butyl, E' is oxygen and (R^c)_q is in the 5-position and corresponds in each case to a row of Table B.

Table 119

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ is 25 hydrogen and R² is tert-butyl, E' is oxygen and (R^c)_q is in the 5-position and corresponds in each case to a row of Table B.

Table 120

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ is 30 hydrogen and R² is phenyl, E' is oxygen and (R^c)_q is in the 5-position and corresponds in each case to a row of Table B.

Table 121

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ and 35 R² are hydrogen, E' is sulfur and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

Table 122

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is 40 hydrogen and R² is methyl, E' is sulfur and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

Table 123

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is 45 hydrogen and R² is ethyl, E' is sulfur and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

Table 124

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is i-propyl, E' is sulfur and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

5

Table 125

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is n-propyl, E' is sulfur and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

10

Table 126

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is cyclo-propyl, E' is sulfur and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

15

Table 127

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is n-butyl, E' is sulfur and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

20

Table 128

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is i-butyl, E' is sulfur and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

25

Table 129

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is tert-butyl, E' is sulfur and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

30

Table 130

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is phenyl, E' is sulfur and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

35

Table 131

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ and R² are hydrogen, E' is sulfur and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

40

Table 132

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is hydrogen and R² is methyl, E' is sulfur and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

45

Table 133

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is

39

hydrogen and R^2 is ethyl, E' is sulfur and $(R^c)_q$ is in the 4-position and corresponds in each case to a row of Table B.

Table 134

- 5 Compounds of the formula I.6 wherein X is hydrogen, p is 4, R^1 is hydrogen and R^2 is i-propyl, E' is sulfur and $(R^c)_q$ is in the 4-position and corresponds in each case to a row of Table B.

Table 135

- 10 Compounds of the formula I.6 wherein X is hydrogen, p is 4, R^1 is hydrogen and R^2 is n-propyl, E' is sulfur and $(R^c)_q$ is in the 4-position and corresponds in each case to a row of Table B.

Table 136

- 15 Compounds of the formula I.6 wherein X is hydrogen, p is 4, R^1 is hydrogen and R^2 is cyclo-propyl, E' is sulfur and $(R^c)_q$ is in the 4-position and corresponds in each case to a row of Table B.

Table 137

- 20 Compounds of the formula I.6 wherein X is hydrogen, p is 4, R^1 is hydrogen and R^2 is n-butyl, E' is sulfur and $(R^c)_q$ is in the 4-position and corresponds in each case to a row of Table B.

Table 138

- 25 Compounds of the formula I.6 wherein X is hydrogen, p is 4, R^1 is hydrogen and R^2 is i-butyl, E' is sulfur and $(R^c)_q$ is in the 4-position and corresponds in each case to a row of Table B.

Table 139

- 30 Compounds of the formula I.6 wherein X is hydrogen, p is 4, R^1 is hydrogen and R^2 is tert-butyl, E' is sulfur and $(R^c)_q$ is in the 4-position and corresponds in each case to a row of Table B.

Table 140

- 35 Compounds of the formula I.6 wherein X is hydrogen, p is 4, R^1 is hydrogen and R^2 is phenyl, E' is sulfur and $(R^c)_q$ is in the 4-position and corresponds in each case to a row of Table B.

Table 141

- 40 Compounds of the formula I.6 wherein X is fluorine, p is 2, R^1 and R^2 are hydrogen, E' is sulfur and $(R^c)_q$ is in the 4-position and corresponds in each case to a row of Table B.

Table 142

- 45 Compounds of the formula I.6 wherein X is fluorine, p is 2, R^1 is hydrogen and R^2 is methyl, E' is sulfur and $(R^c)_q$ is in the

40

4-position and corresponds in each case to a row of Table B.

Table 143

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is 5 hydrogen and R² is ethyl, E' is sulfur and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

Table 144

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is 10 hydrogen and R² is i-propyl, E' is sulfur and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

Table 145

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is 15 hydrogen and R² is n-propyl, E' is sulfur and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

Table 146

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is 20 hydrogen and R² is cyclo-propyl, E' is sulfur and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

Table 147

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is 25 hydrogen and R² is n-butyl, E' is sulfur and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

Table 148

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is 30 hydrogen and R² is i-butyl, E' is sulfur and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

Table 149

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is 35 hydrogen and R² is tert-butyl, E' is sulfur and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

Table 150

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is 40 hydrogen and R² is phenyl, E' is sulfur and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

Table 151

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ and 45 R² are hydrogen, E' is sulfur and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

41

Table 152

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ is hydrogen and R² is methyl, E' is sulfur and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

5

Table 153

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ is hydrogen and R² is ethyl, E' is sulfur and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

10

Table 154

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ is hydrogen and R² is i-propyl, E' is sulfur and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

15

Table 155

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ is hydrogen and R² is n-propyl, E' is sulfur and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

20

Table 156

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ is hydrogen and R² is cyclo-propyl, E' is sulfur and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

25

Table 157

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ is hydrogen and R² is n-butyl, E' is sulfur and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

30

Table 158

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ is hydrogen and R² is i-butyl, E' is sulfur and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

35

Table 159

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ is hydrogen and R² is tert-butyl, E' is sulfur and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

40

Table 160

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ is hydrogen and R² is phenyl, E' is sulfur and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

45

Table 161

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ and

42

R² are hydrogen, E' is sulfur and (R^c)_q is in the 5-position and corresponds in each case to a row of Table B.

Table 162

- 5 Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is methyl, E' is sulfur and (R^c)_q is in the 5-position and corresponds in each case to a row of Table B.

Table 163

- 10 Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is ethyl, E' is sulfur and (R^c)_q is in the 5-position and corresponds in each case to a row of Table B.

Table 164

- 15 Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is i-propyl, E' is sulfur and (R^c)_q is in the 5-position and corresponds in each case to a row of Table B.

Table 165

- 20 Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is n-propyl, E' is sulfur and (R^c)_q is in the 5-position and corresponds in each case to a row of Table B.

Table 166

- 25 Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is cyclo-propyl, E' is sulfur and (R^c)_q is in the 5-position and corresponds in each case to a row of Table B.

Table 167

- 30 Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is n-butyl, E' is sulfur and (R^c)_q is in the 5-position and corresponds in each case to a row of Table B.

Table 168

- 35 Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is i-butyl, E' is sulfur and (R^c)_q is in the 5-position and corresponds in each case to a row of Table B.

Table 169

- 40 Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is tert-butyl, E' is sulfur and (R^c)_q is in the 5-position and corresponds in each case to a row of Table B.

Table 170

- 45 Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is phenyl, E' is sulfur and (R^c)_q is in the

43

5-position and corresponds in each case to a row of Table B.

Table 171

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ and
5 R² are hydrogen, E' is sulfur and (R^c)_q is in the 5-position and
corresponds in each case to a row of Table B.

Table 172

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is
10 hydrogen and R² is methyl, E' is sulfur and (R^c)_q is in the
5-position and corresponds in each case to a row of Table B.

Table 173

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is
15 hydrogen and R² is ethyl, E' is sulfur and (R^c)_q is in the
5-position and corresponds in each case to a row of Table B.

Table 174

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is
20 hydrogen and R² is i-propyl, E' is sulfur and (R^c)_q is in the
5-position and corresponds in each case to a row of Table B.

Table 175

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is
25 hydrogen and R² is n-propyl, E' is sulfur and (R^c)_q is in the
5-position and corresponds in each case to a row of Table B.

Table 176

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is
30 hydrogen and R² is cyclo-propyl, E' is sulfur and (R^c)_q is in the
5-position and corresponds in each case to a row of Table B.

Table 177

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is
35 hydrogen and R² is n-butyl, E' is sulfur and (R^c)_q is in the
5-position and corresponds in each case to a row of Table B.

Table 178

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is
40 hydrogen and R² is i-butyl, E' is sulfur and (R^c)_q is in the
5-position and corresponds in each case to a row of Table B.

Table 179

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is
45 hydrogen and R² is tert-butyl, E' is sulfur and (R^c)_q is in the
5-position and corresponds in each case to a row of Table B.

Table 180

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is hydrogen and R² is phenyl, E' is sulfur and (R^c)_q is in the 5-position and corresponds in each case to a row of Table B.

5

Table 181

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ and R² are hydrogen, E' is sulfur and (R^c)_q is in the 5-position and corresponds in each case to a row of Table B.

10

Table 182

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is hydrogen and R² is methyl, E' is sulfur and (R^c)_q is in the 5-position and corresponds in each case to a row of Table B.

15

Table 183

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is hydrogen and R² is ethyl, E' is sulfur and R^c is in the 5-position and corresponds in each case to a row of Table B.

20

Table 184

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is hydrogen and R² is i-propyl, E' is sulfur and (R^c)_q is in the 5-position and corresponds in each case to a row of Table B.

25

Table 185

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is hydrogen and R² is n-propyl, E' is sulfur and (R^c)_q is in the 5-position and corresponds in each case to a row of Table B.

30

Table 186

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is hydrogen and R² is cyclo-propyl, E' is sulfur and (R^c)_q is in the 5-position and corresponds in each case to a row of Table B.

35

Table 187

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is hydrogen and R² is n-butyl, E' is sulfur and (R^c)_q is in the 5-position and corresponds in each case to a row of Table B.

40

Table 188

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is hydrogen and R² is i-butyl, E' is sulfur and (R^c)_q is in the 5-position and corresponds in each case to a row of Table B.

45

Table 189

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is

45

hydrogen and R^2 is tert-butyl, E' is sulfur and $(R^c)_q$ is in the 5-position and corresponds in each case to a row of Table B.

Table 190

- 5 Compounds of the formula I.6 wherein X is fluorine, p is 2, R^1 is hydrogen and R^2 is phenyl, E' is sulfur and $(R^c)_q$ is in the 5-position and corresponds in each case to a row of Table B.

Table 191

- 10 Compounds of the formula I.6 wherein X is fluorine, p is 4, R^1 and R^2 are hydrogen, E' is sulfur and $(R^c)_q$ is in the 5-position and corresponds in each case to a row of Table B.

Table 192

- 15 Compounds of the formula I.6 wherein X is fluorine, p is 4, R^1 is hydrogen and R^2 is methyl, E' is sulfur and $(R^c)_q$ is in the 5-position and corresponds in each case to a row of Table B.

Table 193

- 20 Compounds of the formula I.6 wherein X is fluorine, p is 4, R^1 is hydrogen and R^2 is ethyl, E' is sulfur and $(R^c)_q$ is in the 5-position and corresponds in each case to a row of Table B.

Table 194

- 25 Compounds of the formula I.6 wherein X is fluorine, p is 4, R^1 is hydrogen and R^2 is i-propyl, E' is sulfur and $(R^c)_q$ is in the 5-position and corresponds in each case to a row of Table B.

Table 195

- 30 Compounds of the formula I.6 wherein X is fluorine, p is 4, R^1 is hydrogen and R^2 is n-propyl, E' is sulfur and $(R^c)_q$ is in the 5-position and corresponds in each case to a row of Table B.

Table 196

- 35 Compounds of the formula I.6 wherein X is fluorine, p is 4, R^1 is hydrogen and R^2 is cyclo-propyl, E' is sulfur and $(R^c)_q$ is in the 5-position and corresponds in each case to a row of Table B.

Table 197

- 40 Compounds of the formula I.6 wherein X is fluorine, p is 4, R^1 is hydrogen and R^2 is n-butyl, E' is sulfur and $(R^c)_q$ is in the 5-position and corresponds in each case to a row of Table B.

Table 198

- 45 Compounds of the formula I.6 wherein X is fluorine, p is 4, R^1 is hydrogen and R^2 is i-butyl, E' is sulfur and $(R^c)_q$ is in the

5-position and corresponds in each case to a row of Table B.

Table 199

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ is
5 hydrogen and R² is tert-butyl, E' is sulfur and (R^c)_q is in the
5-position and corresponds in each case to a row of Table B.

Table 200

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ is
10 hydrogen and R² is phenyl, E' is sulfur and (R^c)_q is in the
5-position and corresponds in each case to a row of Table B.

Table 201

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ and
15 R² are hydrogen, E' is NH and (R^c)_q is in the 4-position and
corresponds in each case to a row of Table B.

Table 202

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is
20 hydrogen and R² is methyl, E' is NH and (R^c)_q is in the 4-position
and corresponds in each case to a row of Table B.

Table 203

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is
25 hydrogen and R² is ethyl, E' is NH and (R^c)_q is in the 4-position
and corresponds in each case to a row of Table B.

Table 204

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is
30 hydrogen and R² is i-propyl, E' is NH and (R^c)_q is in the
4-position and corresponds in each case to a row of Table B.

Table 205

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is
35 hydrogen and R² is n-propyl, E' is NH and (R^c)_q is in the
4-position and corresponds in each case to a row of Table B.

Table 206

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is
40 hydrogen and R² is cyclo-propyl, E' is NH and (R^c)_q is in the
4-position and corresponds in each case to a row of Table B.

Table 207

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is
45 hydrogen and R² is n-butyl, E' is NH and (R^c)_q is in the
4-position and corresponds in each case to a row of Table B.

Table 208

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is i-butyl, E' is NH and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

5

Table 209

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is tert-butyl, E' is NH and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

10

Table 210

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is phenyl, E' is NH and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

15

Table 211

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ and R² are hydrogen, E' is NH and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

20

Table 212

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is hydrogen and R² is methyl, E' is NH and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

25

Table 213

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is hydrogen and R² is ethyl, E' is NH and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

30

Table 214

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is hydrogen and R² is i-propyl, E' is NH and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

35

Table 215

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is hydrogen and R² is n-propyl, E' is NH and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

40

Table 216

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is hydrogen and R² is cyclo-propyl, E' is NH and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

45

Table 217

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is

48

hydrogen and R^2 is n-butyl, E' is NH and $(R^c)_q$ is in the 4-position and corresponds in each case to a row of Table B.

Table 218

- 5 Compounds of the formula I.6 wherein X is hydrogen, p is 4, R^1 is hydrogen and R^2 is i-butyl, E' is NH and $(R^c)_q$ is in the 4-position and corresponds in each case to a row of Table B.

Table 219

- 10 Compounds of the formula I.6 wherein X is hydrogen, p is 4, R^1 is hydrogen and R^2 is tert-butyl, E' is NH and $(R^c)_q$ is in the 4-position and corresponds in each case to a row of Table B.

Table 220

- 15 Compounds of the formula I.6 wherein X is hydrogen, p is 4, R^1 is hydrogen and R^2 is phenyl, E' is NH and $(R^c)_q$ is in the 4-position and corresponds in each case to a row of Table B.

Table 221

- 20 Compounds of the formula I.6 wherein X is fluorine, p is 2, R^1 and R^2 are hydrogen, E' is NH and $(R^c)_q$ is in the 4-position and corresponds in each case to a row of Table B.

Table 222

- 25 Compounds of the formula I.6 wherein X is fluorine, p is 2, R^1 is hydrogen and R^2 is methyl, E' is NH and $(R^c)_q$ is in the 4-position and corresponds in each case to a row of Table B.

Table 223

- 30 Compounds of the formula I.6 wherein X is fluorine, p is 2, R^1 is hydrogen and R^2 is ethyl, E' is NH and $(R^c)_q$ is in the 4-position and corresponds in each case to a row of Table B.

Table 224

- 35 Compounds of the formula I.6 wherein X is fluorine, p is 2, R^1 is hydrogen and R^2 is i-propyl, E' is NH and $(R^c)_q$ is in the 4-position and corresponds in each case to a row of Table B.

Table 225

- 40 Compounds of the formula I.6 wherein X is fluorine, p is 2, R^1 is hydrogen and R^2 is n-propyl, E' is NH and $(R^c)_q$ is in the 4-position and corresponds in each case to a row of Table B.

Table 226

- 45 Compounds of the formula I.6 wherein X is fluorine, p is 2, R^1 is hydrogen and R^2 is cyclo-propyl, E' is NH and $(R^c)_q$ is in the

49

4-position and corresponds in each case to a row of Table B.

Table 227

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is
5 hydrogen and R² is n-butyl, E' is NH and (R^c)_q is in the
4-position and corresponds in each case to a row of Table B.

Table 228

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is
10 hydrogen and R² is i-butyl, E' is NH and (R^c)_q is in the
4-position and corresponds in each case to a row of Table B.

Table 229

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is
15 hydrogen and R² is tert-butyl, E' is NH and (R^c)_q is in the
4-position and corresponds in each case to a row of Table B.

Table 230

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is
20 hydrogen and R² is phenyl, E' is NH and (R^c)_q is in the 4-position
and corresponds in each case to a row of Table B.

Table 231

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ and
25 R² are hydrogen, E' is NH and (R^c)_q is in the 4-position and
corresponds in each case to a row of Table B.

Table 232

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ is
30 hydrogen and R² is methyl, E' is NH and (R^c)_q is in the 4-position
and corresponds in each case to a row of Table B.

Table 233

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ is
35 hydrogen and R² is ethyl, E' is NH and (R^c)_q is in the 4-position
and corresponds in each case to a row of Table B.

Table 234

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ is
40 hydrogen and R² is i-propyl, E' is NH and (R^c)_q is in the
4-position and corresponds in each case to a row of Table B.

Table 235

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ is
45 hydrogen and R² is n-propyl, E' is NH and (R^c)_q is in the
4-position and corresponds in each case to a row of Table B.

50

Table 236

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ is hydrogen and R² is cyclo-propyl, E' is NH and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

5

Table 237

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ is hydrogen and R² is n-butyl, E' is NH and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

10

Table 238

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ is hydrogen and R² is i-butyl, E' is NH and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

15

Table 239

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ is hydrogen and R² is tert-butyl, E' is NH and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

20

Table 240

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ is hydrogen and R² is phenyl, E' is NH and (R^c)_q is in the 4-position and corresponds in each case to a row of Table B.

25

Table 241

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ and R² are hydrogen, E' is NH and (R^c)_q is in the 5-position and corresponds in each case to a row of Table B.

30

Table 242

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is methyl, E' is NH and (R^c)_q is in the 5-position and corresponds in each case to a row of Table B.

35

Table 243

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is ethyl, E' is NH and (R^c)_q is in the 5-position and corresponds in each case to a row of Table B.

40

Table 244

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is i-propyl, E' is NH and (R^c)_q is in the 5-position and corresponds in each case to a row of Table B.

45

Table 245

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is

51

hydrogen and R^2 is n-propyl, E' is NH and $(R^c)_q$ is in the 5-position and corresponds in each case to a row of Table B.

Table 246

- 5 Compounds of the formula I.6 wherein X is hydrogen, p is 2, R^1 is hydrogen and R^2 is cyclo-propyl, E' is NH and $(R^c)_q$ is in the 5-position and corresponds in each case to a row of Table B.

Table 247

- 10 Compounds of the formula I.6 wherein X is hydrogen, p is 2, R^1 is hydrogen and R^2 is n-butyl, E' is NH and $(R^c)_q$ is in the 5-position and corresponds in each case to a row of Table B.

Table 248

- 15 Compounds of the formula I.6 wherein X is hydrogen, p is 2, R^1 is hydrogen and R^2 is i-butyl, E' is NH and $(R^c)_q$ is in the 5-position and corresponds in each case to a row of Table B.

Table 249

- 20 Compounds of the formula I.6 wherein X is hydrogen, p is 2, R^1 is hydrogen and R^2 is tert-butyl, E' is NH and $(R^c)_q$ is in the 5-position and corresponds in each case to a row of Table B.

Table 250

- 25 Compounds of the formula I.6 wherein X is hydrogen, p is 2, R^1 is hydrogen and R^2 is phenyl, E' is NH and $(R^c)_q$ is in the 5-position and corresponds in each case to a row of Table B.

Table 251

- 30 Compounds of the formula I.6 wherein X is hydrogen, p is 4, R^1 and R^2 are hydrogen, E' is NH and $(R^c)_q$ is in the 5-position and corresponds in each case to a row of Table B.

Table 252

- 35 Compounds of the formula I.6 wherein X is hydrogen, p is 4, R^1 is hydrogen and R^2 is methyl, E' is NH and $(R^c)_q$ is in the 5-position and corresponds in each case to a row of Table B.

Table 253

- 40 Compounds of the formula I.6 wherein X is hydrogen, p is 4, R^1 is hydrogen and R^2 is ethyl, E' is NH and $(R^c)_q$ is in the 5-position and corresponds in each case to a row of Table B.

Table 254

- 45 Compounds of the formula I.6 wherein X is hydrogen, p is 4, R^1 is hydrogen and R^2 is i-propyl, E' is NH and $(R^c)_q$ is in the

52

5-position and corresponds in each case to a row of Table B.

Table 255

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is
5 hydrogen and R² is n-propyl, E' is NH and (R^c)_q is in the
5-position and corresponds in each case to a row of Table B.

Table 256

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is
10 hydrogen and R² is cyclo-propyl, E' is NH and (R^c)_q is in the
5-position and corresponds in each case to a row of Table B.

Table 257

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is
15 hydrogen and R² is n-butyl, E' is NH and (R^c)_q is in the
5-position and corresponds in each case to a row of Table B.

Table 258

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is
20 hydrogen and R² is i-butyl, E' is NH and (R^c)_q is in the
5-position and corresponds in each case to a row of Table B.

Table 259

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is
25 hydrogen and R² is tert-butyl, E' is NH and (R^c)_q is in the
5-position and corresponds in each case to a row of Table B.

Table 260

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is
30 hydrogen and R² is phenyl, E' is NH and (R^c)_q is in the 5-position
and corresponds in each case to a row of Table B.

Table 261

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ and
35 R² are hydrogen, E' is NH and (R^c)_q is in the 5-position and
corresponds in each case to a row of Table B.

Table 262

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is
40 hydrogen and R² is methyl, E' is NH and (R^c)_q is in the 5-position
and corresponds in each case to a row of Table B.

Table 263

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is
45 hydrogen and R² is ethyl, E' is NH and (R^c)_q is in the 5-position
and corresponds in each case to a row of Table B.

Table 264

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is hydrogen and R² is i-propyl, E' is NH and (R^c)_q is in the 5-position and corresponds in each case to a row of Table B.

5

Table 265

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is hydrogen and R² is n-propyl, E' is NH and (R^c)_q is in the 5-position and corresponds in each case to a row of Table B.

10

Table 266

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is hydrogen and R² is cyclo-propyl, E' is NH and (R^c)_q is in the 5-position and corresponds in each case to a row of Table B.

15

Table 267

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is hydrogen and R² is n-butyl, E' is NH and (R^c)_q is in the 5-position and corresponds in each case to a row of Table B.

20

Table 268

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is hydrogen and R² is i-butyl, E' is NH and (R^c)_q is in the 5-position and corresponds in each case to a row of Table B.

25

Table 269

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is hydrogen and R² is tert-butyl, E' is NH and (R^c)_q is in the 5-position and corresponds in each case to a row of Table B.

30

Table 270

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is hydrogen and R² is phenyl, E' is NH and (R^c)_q is in the 5-position and corresponds in each case to a row of Table B.

35

Table 271

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ and R² are hydrogen, E' is NH and (R^c)_q is in the 5-position and corresponds in each case to a row of Table B.

40

Table 272

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ is hydrogen and R² is methyl, E' is NH and (R^c)_q is in the 5-position and corresponds in each case to a row of Table B.

45

Table 273

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ is

54

hydrogen and R^2 is ethyl, E' is NH and $(R^c)_q$ is in the 5-position and corresponds in each case to a row of Table B.

Table 274

- 5 Compounds of the formula I.6 wherein X is fluorine, p is 4, R^1 is hydrogen and R^2 is i-propyl, E' is NH and $(R^c)_q$ is in the 5-position and corresponds in each case to a row of Table B.

Table 275

- 10 Compounds of the formula I.6 wherein X is fluorine, p is 4, R^1 is hydrogen and R^2 is n-propyl, E' is NH and $(R^c)_q$ is in the 5-position and corresponds in each case to a row of Table B.

Table 276

- 15 Compounds of the formula I.6 wherein X is fluorine, p is 4, R^1 is hydrogen and R^2 is cyclo-propyl, E' is NH and $(R^c)_q$ is in the 5-position and corresponds in each case to a row of Table B.

Table 277

- 20 Compounds of the formula I.6 wherein X is fluorine, p is 4, R^1 is hydrogen and R^2 is n-butyl, E' is NH and R^c is in the 5-position and corresponds in each case to a row of Table B.

Table 278

- 25 Compounds of the formula I.6 wherein X is fluorine, p is 4, R^1 is hydrogen and R^2 is i-butyl, E' is NH and $(R^c)_q$ is in the 5-position and corresponds in each case to a row of Table B.

Table 279

- 30 Compounds of the formula I.6 wherein X is fluorine, p is 4, R^1 is hydrogen and R^2 is tert-butyl, E' is NH and $(R^c)_q$ is in the 5-position and corresponds in each case to a row of Table B.

Table 280

- 35 Compounds of the formula I.6 wherein X is fluorine, p is 4, R^1 is hydrogen and R^2 is phenyl, E' is NH and $(R^c)_q$ is in the 5-position and corresponds in each case to a row of Table B.




Table B

45

Nr.	$(R^c)_q$
B-1	-
B-2	CH ₃

55

Nr.	$(R^C)_q$
B-3	CH_2CH_3
B-4	$CH_2CH_2CH_3$
5	B-5 $CH(CH_3)_2$
B-6	$C(CH_3)_3$
B-7	Cl
B-8	Br
10	B-9 CH_2OCH_3
B-10	$COOCH_3$
B-11	C_6H_5
B-12	$2-F-C_6H_4$
B-13	$3-F-C_6H_4$
15	B-14 $4-F-C_6H_4$
B-15	$2-Cl-C_6H_4$
B-16	$3-Cl-C_6H_4$
B-17	$4-Cl-C_6H_4$
20	B-18 $2-CH_3-C_6H_4$
B-19	$3-CH_3-C_6H_4$
B-20	$4-CH_3-C_6H_4$
B-21	$2,3-(CH_3)_2-C_6H_3$
25	B-22 $2,4-(CH_3)_2-C_6H_3$
B-23	$2,5-(CH_3)_2-C_6H_3$
B-24	$3,4-(CH_3)_2-C_6H_3$
B-25	$3,5-(CH_3)_2-C_6H_3$
30	B-26 $2,3-Cl_2-C_6H_3$
B-27	$2,4-Cl_2-C_6H_3$
B-28	$2,5-Cl_2-C_6H_3$
B-29	$3,4-Cl_2-C_6H_3$
B-30	$3,5-Cl_2-C_6H_3$
35	B-31 $2,3-F_2-C_6H_3$
B-32	$2,4-F_2-C_6H_3$
B-33	$2,5-F_2-C_6H_3$
B-34	$3,4-F_2-C_6H_3$
40	B-35 $3,5-F_2-C_6H_3$
B-36	$2-F-3-Cl-C_6H_3$
B-37	$2-F-4-Cl-C_6H_3$
B-38	$2-F-5-Cl-C_6H_3$
45	B-39 $3-F-4-Cl-C_6H_3$
B-40	$3-F-5-Cl-C_6H_3$
B-41	$2-Cl-3-F-C_6H_3$

		Nr.	(R ^c) _q
5		B-42	2-Cl-4-F-C ₆ H ₃
		B-43	2-Cl-5-F-C ₆ H ₃
		B-44	3-Cl-4-F-C ₆ H ₃
		B-45	3-Cl-5-F-C ₆ H ₃
		B-46	2-F-3-CH ₃ -C ₆ H ₃
10		B-47	2-F-4-CH ₃ -C ₆ H ₃
		B-48	2-F-5-CH ₃ -C ₆ H ₃
		B-49	3-F-4-CH ₃ -C ₆ H ₃
		B-50	3-F-5-CH ₃ -C ₆ H ₃
		B-51	2-CH ₃ -3-F-C ₆ H ₃
15		B-52	2-CH ₃ -4-F-C ₆ H ₃
		B-53	2-CH ₃ -5-F-C ₆ H ₃
		B-54	3-CH ₃ -4-F-C ₆ H ₃
		B-55	3-CH ₃ -5-F-C ₆ H ₃
		B-56	2-Cl-3-CH ₃ -C ₆ H ₃
20		B-57	2-Cl-4-CH ₃ -C ₆ H ₃
		B-58	2-Cl-5-CH ₃ -C ₆ H ₃
		B-59	3-Cl-4-CH ₃ -C ₆ H ₃
		B-60	3-Cl-5-CH ₃ -C ₆ H ₃
		B-61	2-CH ₃ -3-Cl-C ₆ H ₃
25		B-62	2-CH ₃ -4-Cl-C ₆ H ₃
		B-63	2-CH ₃ -5-Cl-C ₆ H ₃
		B-64	3-CH ₃ -4-Cl-C ₆ H ₃
		B-65	3-CH ₃ -5-Cl-C ₆ H ₃
		B-66	2,3,4-(CH ₃) ₃ -C ₆ H ₂
30		B-67	2,3,5-(CH ₃) ₃ -C ₆ H ₂
		B-68	2,3,6-(CH ₃) ₃ -C ₆ H ₂
		B-69	2,4,5-(CH ₃) ₃ -C ₆ H ₂
		B-70	2,4,6-(CH ₃) ₃ -C ₆ H ₂
		B-71	3,4,5-(CH ₃) ₃ -C ₆ H ₂
35		B-72	3,4,6-(CH ₃) ₃ -C ₆ H ₂
		B-73	2,3,4-F ₃ -C ₆ H ₂
		B-74	2,3,5-F ₃ -C ₆ H ₂
		B-75	2,3,6-F ₃ -C ₆ H ₂
		B-76	2,4,5-F ₃ -C ₆ H ₂
40		B-77	2,4,6-F ₃ -C ₆ H ₂
		B-78	3,4,5-F ₃ -C ₆ H ₂
		B-79	3,4,6-F ₃ -C ₆ H ₂
		B-80	2,3,4-Cl ₃ -C ₆ H ₂

57

Nr.	$(R^c)_q$
5	B-81 2,3,5-Cl ₃ -C ₆ H ₂
	B-82 2,3,6-Cl ₃ -C ₆ H ₂
	B-83 2,4,5-Cl ₃ -C ₆ H ₂
	B-84 2,4,6-Cl ₃ -C ₆ H ₂
	B-85 3,4,5-Cl ₃ -C ₆ H ₂
	B-86 3,4,6-Cl ₃ -C ₆ H ₂
10	B-87 2-Cl-3,4-(CH ₃) ₂ -C ₆ H ₂
	B-88 3-Cl-2,4-(CH ₃) ₂ -C ₆ H ₂
	B-89 4-Cl-2,3-(CH ₃) ₂ -C ₆ H ₂
	B-90 2-Cl-3,5-(CH ₃) ₂ -C ₆ H ₂
	B-91 3-Cl-2,5-(CH ₃) ₂ -C ₆ H ₂
15	B-92 5-Cl-2,3-(CH ₃) ₂ -C ₆ H ₂
	B-93 2-Cl-3,6-(CH ₃) ₂ -C ₆ H ₂
	B-94 3-Cl-2,6-(CH ₃) ₂ -C ₆ H ₂
	B-95 6-Cl-2,3-(CH ₃) ₂ -C ₆ H ₂
20	B-96 2-Cl-4,5-(CH ₃) ₂ -C ₆ H ₂
	B-97 4-Cl-2,5-(CH ₃) ₂ -C ₆ H ₂
	B-98 5-Cl-2,4-(CH ₃) ₂ -C ₆ H ₂
	B-99 2-Cl-4,6-(CH ₃) ₂ -C ₆ H ₂
25	B-100 4-Cl-2,6-(CH ₃) ₂ -C ₆ H ₂
	B-101 6-Cl-2,4-(CH ₃) ₂ -C ₆ H ₂
	B-102 3-Cl-4,5-(CH ₃) ₂ -C ₆ H ₂
	B-103 4-Cl-3,5-(CH ₃) ₂ -C ₆ H ₂
	B-104 5-Cl-3,4-(CH ₃) ₂ -C ₆ H ₂
30	B-105 3-Cl-4,6-(CH ₃) ₂ -C ₆ H ₂
	B-106 4-Cl-3,6-(CH ₃) ₂ -C ₆ H ₂
	B-107 6-Cl-3,4-(CH ₃) ₂ -C ₆ H ₂
	B-108 3-Cl-5,6-(CH ₃) ₂ -C ₆ H ₂
35	B-109 5-Cl-3,6-(CH ₃) ₂ -C ₆ H ₂
	B-110 6-Cl-3,5-(CH ₃) ₂ -C ₆ H ₂
	B-111 2-F-3,4-(CH ₃) ₂ -C ₆ H ₂
	B-112 3-F-2,4-(CH ₃) ₂ -C ₆ H ₂
40	B-113 4-F-2,3-(CH ₃) ₂ -C ₆ H ₂
	B-114 2-F-3,5-(CH ₃) ₂ -C ₆ H ₂
	B-115 3-F-2,5-(CH ₃) ₂ -C ₆ H ₂
	B-116 5-F-2,3-(CH ₃) ₂ -C ₆ H ₂
45	B-117 2-F-3,6-(CH ₃) ₂ -C ₆ H ₂
	B-118 3-F-2,6-(CH ₃) ₂ -C ₆ H ₂
	B-119 6-F-2,3-(CH ₃) ₂ -C ₆ H ₂

		Nr.	(R ^a) _q
5		B-120	2-F-4, 5-(CH ₃) ₂ -C ₆ H ₂
		B-121	4-F-2, 5-(CH ₃) ₂ -C ₆ H ₂
		B-122	5-F-2, 4-(CH ₃) ₂ -C ₆ H ₂
		B-123	2-F-4, 6-(CH ₃) ₂ -C ₆ H ₂
		B-124	4-F-2, 6-(CH ₃) ₂ -C ₆ H ₂
10		B-125	6-F-2, 4-(CH ₃) ₂ -C ₆ H ₂
		B-126	3-F-4, 5-(CH ₃) ₂ -C ₆ H ₂
		B-127	4-F-3, 5-(CH ₃) ₂ -C ₆ H ₂
		B-128	5-F-3, 4-(CH ₃) ₂ -C ₆ H ₂
		B-129	3-F-4, 6-(CH ₃) ₂ -C ₆ H ₂
15		B-130	4-F-3, 6-(CH ₃) ₂ -C ₆ H ₂
		B-131	6-F-3, 4-(CH ₃) ₂ -C ₆ H ₂
		B-132	3-F-5, 6-(CH ₃) ₂ -C ₆ H ₂
		B-133	5-F-3, 6-(CH ₃) ₂ -C ₆ H ₂
		B-134	6-F-3, 5-(CH ₃) ₂ -C ₆ H ₂
20		B-135	2-Cl-3, 4-F ₂ -C ₆ H ₂
		B-136	3-Cl-2, 4-F ₂ -C ₆ H ₂
		B-137	4-Cl-2, 3-F ₂ -C ₆ H ₂
		B-138	2-Cl-3, 5-F ₂ -C ₆ H ₂
		B-139	3-Cl-2, 5-F ₂ -C ₆ H ₂
25		B-140	5-Cl-2, 3-F ₂ -C ₆ H ₂
		B-141	2-Cl-3, 6-F ₂ -C ₆ H ₂
		B-142	3-Cl-2, 6-F ₂ -C ₆ H ₂
		B-143	6-Cl-2, 3-F ₂ -C ₆ H ₂
		B-144	2-Cl-4, 5-F ₂ -C ₆ H ₂
30		B-145	4-Cl-2, 5-F ₂ -C ₆ H ₂
		B-146	5-Cl-2, 4-F ₂ -C ₆ H ₂
		B-147	2-Cl-4, 6-F ₂ -C ₆ H ₂
		B-148	4-Cl-2, 6-F ₂ -C ₆ H ₂
		B-149	6-Cl-2, 4-F ₂ -C ₆ H ₂
35		B-150	3-Cl-4, 5-F ₂ -C ₆ H ₂
		B-151	4-Cl-3, 5-F ₂ -C ₆ H ₂
		B-152	5-Cl-3, 4-F ₂ -C ₆ H ₂
		B-153	3-Cl-4, 6-F ₂ -C ₆ H ₂
		B-154	4-Cl-3, 6-F ₂ -C ₆ H ₂
40		B-155	6-Cl-3, 4-F ₂ -C ₆ H ₂
		B-156	3-Cl-5, 6-F ₂ -C ₆ H ₂
		B-157	5-Cl-3, 6-F ₂ -C ₆ H ₂
		B-158	6-Cl-3, 5-F ₂ -C ₆ H ₂


		Nr.	(R ^c) _q
5		B-159	2-F-3,4-Cl ₂ -C ₆ H ₂
		B-160	3-F-2,4-Cl ₂ -C ₆ H ₂
		B-161	4-F-2,3-Cl ₂ -C ₆ H ₂
		B-162	2-F-3,5-Cl ₂ -C ₆ H ₂
		B-163	3-F-2,5-Cl ₂ -C ₆ H ₂
10		B-164	5-F-2,3-Cl ₂ -C ₆ H ₂
		B-165	2-F-3,6-Cl ₂ -C ₆ H ₂
		B-166	3-F-2,6-Cl ₂ -C ₆ H ₂
		B-167	6-F-2,3-Cl ₂ -C ₆ H ₂
		B-168	2-F-4,5-Cl ₂ -C ₆ H ₂
15		B-169	4-F-2,5-Cl ₂ -C ₆ H ₂
		B-170	5-F-2,4-Cl ₂ -C ₆ H ₂
		B-171	2-F-4,6-Cl ₂ -C ₆ H ₂
		B-172	4-F-2,6-Cl ₂ -C ₆ H ₂
		B-173	6-F-2,4-Cl ₂ -C ₆ H ₂
20		B-174	3-F-4,5-Cl ₂ -C ₆ H ₂
		B-175	4-F-3,5-Cl ₂ -C ₆ H ₂
		B-176	5-F-3,4-Cl ₂ -C ₆ H ₂
		B-177	3-F-4,6-Cl ₂ -C ₆ H ₂
		B-178	4-F-3,6-Cl ₂ -C ₆ H ₂
25		B-179	6-F-3,4-Cl ₂ -C ₆ H ₂
		B-180	3-F-5,6-Cl ₂ -C ₆ H ₂
		B-181	5-F-3,6-Cl ₂ -C ₆ H ₂
		B-182	6-F-3,5-Cl ₂ -C ₆ H ₂
		B-183	2-CH ₃ -3,4-F ₂ -C ₆ H ₂
30		B-184	3-CH ₃ -2,4-F ₂ -C ₆ H ₂
		B-185	4-CH ₃ -2,3-F ₂ -C ₆ H ₂
		B-186	2-CH ₃ -3,5-F ₂ -C ₆ H ₂
		B-187	3-CH ₃ -2,5-F ₂ -C ₆ H ₂
		B-188	5-CH ₃ -2,3-F ₂ -C ₆ H ₂
35		B-189	2-CH ₃ -3,6-F ₂ -C ₆ H ₂
		B-190	3-CH ₃ -2,6-F ₂ -C ₆ H ₂
		B-191	6-CH ₃ -2,3-F ₂ -C ₆ H ₂
		B-192	2-CH ₃ -4,5-F ₂ -C ₆ H ₂
		B-193	4-CH ₃ -2,5-F ₂ -C ₆ H ₂
40		B-194	5-CH ₃ -2,4-F ₂ -C ₆ H ₂
		B-195	2-CH ₃ -4,6-F ₂ -C ₆ H ₂
		B-196	4-CH ₃ -2,6-F ₂ -C ₆ H ₂
		B-197	6-CH ₃ -2,4-F ₂ -C ₆ H ₂
		B-198	3-CH ₃ -2,5-F ₂ -C ₆ H ₂

60

Nr.	(R ^c) _q
B-198	3-CH ₃ -4,5-F ₂ -C ₆ H ₂
B-199	4-CH ₃ -3,5-F ₂ -C ₆ H ₂
5 B-200	5-CH ₃ -3,4-F ₂ -C ₆ H ₂
B-201	3-CH ₃ -4,6-F ₂ -C ₆ H ₂
B-202	4-CH ₃ -3,6-F ₂ -C ₆ H ₂
B-203	6-CH ₃ -3,4-F ₂ -C ₆ H ₂
10 B-204	3-CH ₃ -5,6-F ₂ -C ₆ H ₂
B-205	5-CH ₃ -3,6-F ₂ -C ₆ H ₂
B-206	6-CH ₃ -3,5-F ₂ -C ₆ H ₂
B-207	2-CH ₃ -3,4-Cl ₂ -C ₆ H ₂
B-208	3-CH ₃ -2,4-Cl ₂ -C ₆ H ₂
15 B-209	4-CH ₃ -2,3-Cl ₂ -C ₆ H ₂
B-210	2-CH ₃ -3,5-Cl ₂ -C ₆ H ₂
B-211	3-CH ₃ -2,5-Cl ₂ -C ₆ H ₂
B-212	5-CH ₃ -2,3-Cl ₂ -C ₆ H ₂
20 B-213	2-CH ₃ -3,6-Cl ₂ -C ₆ H ₂
B-214	3-CH ₃ -2,6-Cl ₂ -C ₆ H ₂
B-215	6-CH ₃ -2,3-Cl ₂ -C ₆ H ₂
B-216	2-CH ₃ -4,5-Cl ₂ -C ₆ H ₂
25 B-217	4-CH ₃ -2,5-Cl ₂ -C ₆ H ₂
B-218	5-CH ₃ -2,4-Cl ₂ -C ₆ H ₂
B-219	2-CH ₃ -4,6-Cl ₂ -C ₆ H ₂
B-220	4-CH ₃ -2,6-Cl ₂ -C ₆ H ₂
B-221	6-CH ₃ -2,4-Cl ₂ -C ₆ H ₂
30 B-222	3-CH ₃ -4,5-Cl ₂ -C ₆ H ₂
B-223	4-CH ₃ -3,5-Cl ₂ -C ₆ H ₂
B-224	5-CH ₃ -3,4-Cl ₂ -C ₆ H ₂
B-225	3-CH ₃ -4,6-Cl ₂ -C ₆ H ₂
35 B-226	4-CH ₃ -3,6-Cl ₂ -C ₆ H ₂
B-227	6-CH ₃ -3,4-Cl ₂ -C ₆ H ₂
B-228	3-CH ₃ -5,6-Cl ₂ -C ₆ H ₂
B-229	5-CH ₃ -3,6-Cl ₂ -C ₆ H ₂
40 B-230	6-CH ₃ -3,5-Cl ₂ -C ₆ H ₂

Table 281

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ and R² are hydrogen, E' is oxygen, q is 2 and (R^c)_q for a compound
 45 corresponds in each case to a row of Table C.

**5 Table 282**

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is methyl, E' is oxygen, q is 2 and (R^c)_q for a compound corresponds in each case to a row of Table C.

10 Table 283

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is ethyl, E' is oxygen, q is 2 and (R^c)_q for a compound corresponds in each case to a row of Table C.

15 Table 284

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is i-propyl, E' is oxygen, q is 2 and (R^c)_q for a compound corresponds in each case to a row of Table C.

20 Table 285

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is n-propyl, E' is oxygen, q is 2 and (R^c)_q for a compound corresponds in each case to a row of Table C.

25 Table 286

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is cyclo-propyl, E' is oxygen, q is 2 and (R^c)_q for a compound corresponds in each case to a row of Table C.

30 Table 287

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is n-butyl, E' is oxygen, q is 2 and (R^c)_q for a compound corresponds in each case to a row of Table C.

35 Table 288

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is i-butyl, E' is oxygen, q is 2 and (R^c)_q for a compound corresponds in each case to a row of Table C.

40 Table 289

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is tert-butyl, E' is oxygen, q is 2 and (R^c)_q for a compound corresponds in each case to a row of Table C.

45 Table 290

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is phenyl, E' is oxygen, q is 2 and (R^c)_q for a

compound corresponds in each case to a row of Table C.

Table 291

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ and
5 R² are hydrogen, E' is oxygen, q is 2 and (R^c)_q for a compound
corresponds in each case to a row of Table C.

Table 292

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is
10 hydrogen and R² is methyl, E' is oxygen, q is 2 and (R^c)_q for a
compound corresponds in each case to a row of Table C.

Table 293

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is
15 hydrogen and R² is ethyl, E' is oxygen, q is 2 and (R^c)_q for a
compound corresponds in each case to a row of Table C.

Table 294

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is
20 hydrogen and R² is i-propyl, E' is oxygen, q is 2 and (R^c)_q for a
compound corresponds in each case to a row of Table C.

Table 295

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is
25 hydrogen and R² is n-propyl, E' is oxygen, q is 2 and (R^c)_q for a
compound corresponds in each case to a row of Table C.

Table 296

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is
30 hydrogen and R² is cyclo-propyl, E' is oxygen, q is 2 and (R^c)_q
for a compound corresponds in each case to a row of Table C.

Table 297

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is
35 hydrogen and R² is n-butyl, E' is oxygen, q is 2 and (R^c)_q for a
compound corresponds in each case to a row of Table C.

Table 298

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is
40 hydrogen and R² is i-butyl, E' is oxygen, q is 2 and (R^c)_q for a
compound corresponds in each case to a row of Table C.

Table 299

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is
45 hydrogen and R² is tert-butyl, E' is oxygen, q is 2 and (R^c)_q for
a compound corresponds in each case to a row of Table C.

Table 300

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is hydrogen and R² is phenyl, E' is oxygen, q is 2 and (R^c)_q for a compound corresponds in each case to a row of Table C.

5

Table 301

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ and R² are hydrogen, E' is oxygen, q is 2 and (R^c)_q for a compound corresponds in each case to a row of Table C.

10

Table 302

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is hydrogen and R² is methyl, E' is oxygen, q is 2 and (R^c)_q for a compound corresponds in each case to a row of Table C.

15

Table 303

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is hydrogen and R² is ethyl, E' is oxygen, q is 2 and (R^c)_q for a compound corresponds in each case to a row of Table C.

20

Table 304

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is hydrogen and R² is i-propyl, E' is oxygen, q is 2 and (R^c)_q for a compound corresponds in each case to a row of Table C.

25

Table 305

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is hydrogen and R² is n-propyl, E' is oxygen, q is 2 and (R^c)_q for a compound corresponds in each case to a row of Table C.

30

Table 306

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is hydrogen and R² is cyclo-propyl, E' is oxygen, q is 2 and (R^c)_q for a compound corresponds in each case to a row of Table C.

35

Table 307

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is hydrogen and R² is n-butyl, E' is oxygen, q is 2 and (R^c)_q for a compound corresponds in each case to a row of Table C.

40

Table 308

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is hydrogen and R² is i-butyl, E' is oxygen, q is 2 and (R^c)_q for a compound corresponds in each case to a row of Table C.

45

Table 309

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is

64

hydrogen and R^2 is tert-butyl, E' is oxygen, q is 2 and $(R^c)_q$ for a compound corresponds in each case to a row of Table C.

Table 310

- 5 Compounds of the formula I.6 wherein X is fluorine, p is 2, R^1 is hydrogen and R^2 is phenyl, E' is oxygen, q is 2 and $(R^c)_q$ for a compound corresponds in each case to a row of Table C.

Table 311

- 10 Compounds of the formula I.6 wherein X is fluorine, p is 4, R^1 and R^2 are hydrogen, E' is oxygen, q is 2 and $(R^c)_q$ for a compound corresponds in each case to a row of Table C.

Table 312

- 15 Compounds of the formula I.6 wherein X is fluorine, p is 4, R^1 is hydrogen and R^2 is methyl, E' is oxygen, q is 2 and $(R^c)_q$ for a compound corresponds in each case to a row of Table C.

Table 313

- 20 Compounds of the formula I.6 wherein X is fluorine, p is 4, R^1 is hydrogen and R^2 is ethyl, E' is oxygen, q is 2 and $(R^c)_q$ for a compound corresponds in each case to a row of Table C.

Table 314

- 25 Compounds of the formula I.6 wherein X is fluorine, p is 4, R^1 is hydrogen and R^2 is i-propyl, E' is oxygen, q is 2 and $(R^c)_q$ for a compound corresponds in each case to a row of Table C.

Table 315

- 30 Compounds of the formula I.6 wherein X is fluorine, p is 4, R^1 is hydrogen and R^2 is n-propyl, E' is oxygen, q is 2 and $(R^c)_q$ for a compound corresponds in each case to a row of Table C.

Table 316

- 35 Compounds of the formula I.6 wherein X is fluorine, p is 4, R^1 is hydrogen and R^2 is cyclo-propyl, E' is oxygen, q is 2 and $(R^c)_q$ for a compound corresponds in each case to a row of Table C.

Table 317

- 40 Compounds of the formula I.6 wherein X is fluorine, p is 4, R^1 is hydrogen and R^2 is n-butyl, E' is oxygen, q is 2 and $(R^c)_q$ for a compound corresponds in each case to a row of Table C.

Table 318

- 45 Compounds of the formula I.6 wherein X is fluorine, p is 4, R^1 is hydrogen and R^2 is i-butyl, E' is oxygen, q is 2 and $(R^c)_q$ for a

65

compound corresponds in each case to a row of Table C.

Table 319

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ is
5 hydrogen and R² is tert-butyl, E' is oxygen, q is 2 and (R^c)_q for
a compound corresponds in each case to a row of Table C.

Table 320

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ is
10 hydrogen and R² is phenyl, E' is oxygen, q is 2 and (R^c)_q for a
compound corresponds in each case to a row of Table C.

Table 321

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ and
15 R² are hydrogen, E' is sulfur, q is 2 and (R^c)_q for a compound
corresponds in each case to a row of Table C.

Table 322

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is
20 hydrogen and R² is methyl, E' is sulfur and (R^c)_q for a compound
corresponds in each case to a row of Table C.

Table 323

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is
25 hydrogen and R² is ethyl, E' is sulfur, q is 2 and (R^c)_q for a
compound corresponds in each case to a row of Table C.

Table 324

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is
30 hydrogen and R² is i-propyl, E' is sulfur, q is 2 and (R^c)_q for a
compound corresponds in each case to a row of Table C.

Table 325

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is
35 hydrogen and R² is n-propyl, E' is sulfur, q is 2 and (R^c)_q for a
compound corresponds in each case to a row of Table C.

Table 326

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is
40 hydrogen and R² is cyclo-propyl, E' is sulfur, q is 2 and (R^c)_q
for a compound corresponds in each case to a row of Table C.

Table 327

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is
45 hydrogen and R² is n-butyl, E' is sulfur, q is 2 and (R^c)_q for a
compound corresponds in each case to a row of Table C.

Table 328

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is i-butyl, E' is sulfur, q is 2 and (R^c)_q for a compound corresponds in each case to a row of Table C.

5

Table 329

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is tert-butyl, E' is sulfur, q is 2 and (R^c)_q for a compound corresponds in each case to a row of Table C.

10

Table 330

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is phenyl, E' is sulfur, q is 2 and (R^c)_q for a compound corresponds in each case to a row of Table C.

15

Table 331

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ and R² are hydrogen, E' is sulfur, q is 2 and (R^c)_q for a compound corresponds in each case to a row of Table C.

20

Table 332

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is hydrogen and R² is methyl, E' is sulfur, q is 2 and (R^c)_q for a compound corresponds in each case to a row of Table C.

25

Table 333

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is hydrogen and R² is ethyl, E' is sulfur, q is 2 and (R^c)_q for a compound corresponds in each case to a row of Table C.

30

Table 334

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is hydrogen and R² is i-propyl, E' is sulfur, q is 2 and (R^c)_q for a compound corresponds in each case to a row of Table C.

35

Table 335

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is hydrogen and R² is n-propyl, E' is sulfur, q is 2 and (R^c)_q for a compound corresponds in each case to a row of Table C.

40

Table 336

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is hydrogen and R² is cyclo-propyl, E' is sulfur, q is 2 and (R^c)_q for a compound corresponds in each case to a row of Table C.

45

Table 337

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is

67

hydrogen and R^2 is n-butyl, E' is sulfur, q is 2 and $(R^c)_q$ for a compound corresponds in each case to a row of Table C.

Table 338

- 5 Compounds of the formula I.6 wherein X is hydrogen, p is 4, R^1 is hydrogen and R^2 is i-butyl, E' is sulfur, q is 2 and $(R^c)_q$ for a compound corresponds in each case to a row of Table C.

Table 339

- 10 Compounds of the formula I.6 wherein X is hydrogen, p is 4, R^1 is hydrogen and R^2 is tert-butyl, E' is sulfur, q is 2 and $(R^c)_q$ for a compound corresponds in each case to a row of Table C.

Table 340

- 15 Compounds of the formula I.6 wherein X is hydrogen, p is 4, R^1 is hydrogen and R^2 is phenyl, E' is sulfur, q is 2 and $(R^c)_q$ for a compound corresponds in each case to a row of Table C.

Table 341

- 20 Compounds of the formula I.6 wherein X is fluorine, p is 2, R^1 and R^2 are hydrogen, E' is sulfur, q is 2 and $(R^c)_q$ for a compound corresponds in each case to a row of Table C.

Table 342

- 25 Compounds of the formula I.6 wherein X is fluorine, p is 2, R^1 is hydrogen and R^2 is methyl, E' is sulfur, q is 2 and $(R^c)_q$ for a compound corresponds in each case to a row of Table C.

Table 343

- 30 Compounds of the formula I.6 wherein X is fluorine, p is 2, R^1 is hydrogen and R^2 is ethyl, E' is sulfur, q is 2 and $(R^c)_q$ for a compound corresponds in each case to a row of Table C.

Table 344

- 35 Compounds of the formula I.6 wherein X is fluorine, p is 2, R^1 is hydrogen and R^2 is i-propyl, E' is sulfur, q is 2 and $(R^c)_q$ for a compound corresponds in each case to a row of Table C.

Table 345

- 40 Compounds of the formula I.6 wherein X is fluorine, p is 2, R^1 is hydrogen and R^2 is n-propyl, E' is sulfur, q is 2 and $(R^c)_q$ for a compound corresponds in each case to a row of Table C.

Table 346

- 45 Compounds of the formula I.6 wherein X is fluorine, p is 2, R^1 is hydrogen and R^2 is cyclo-propyl, E' is sulfur, q is 2 and $(R^c)_q$

for a compound corresponds in each case to a row of Table C.

Table 347

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is
5 hydrogen and R² is n-butyl, E' is sulfur, q is 2 and (R^c)_q for a
compound corresponds in each case to a row of Table C.

Table 348

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is
10 hydrogen and R² is i-butyl, E' is sulfur, q is 2 and (R^c)_q for a
compound corresponds in each case to a row of Table C.

Table 349

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is
15 hydrogen and R² is tert-butyl, E' is sulfur, q is 2 and (R^c)_q for
a compound corresponds in each case to a row of Table C.

Table 350

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is
20 hydrogen and R² is phenyl, E' is sulfur, q is 2 and (R^c)_q for a
compound corresponds in each case to a row of Table C.

Table 351

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ and
25 R² are hydrogen, E' is sulfur, q is 2 and (R^c)_q for a compound
corresponds in each case to a row of Table C.

Table 352

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ is
30 hydrogen and R² is methyl, E' is sulfur, q is 2 and (R^c)_q for a
compound corresponds in each case to a row of Table C.

Table 353

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ is
35 hydrogen and R² is ethyl, E' is sulfur, q is 2 and (R^c)_q for a
compound corresponds in each case to a row of Table C.

Table 354

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ is
40 hydrogen and R² is i-propyl, E' is sulfur, q is 2 and (R^c)_q for a
compound corresponds in each case to a row of Table C.

Table 355

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ is
45 hydrogen and R² is n-propyl, E' is sulfur, q is 2 and (R^c)_q for a
compound corresponds in each case to a row of Table C.

Table 356

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ is hydrogen and R² is cyclo-propyl, E' is sulfur, q is 2 and (R^c)_q for a compound corresponds in each case to a row of Table C.

5

Table 357

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ is hydrogen and R² is n-butyl, E' is sulfur, q is 2 and (R^c)_q for a compound corresponds in each case to a row of Table C.

10

Table 358

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ is hydrogen and R² is i-butyl, E' is sulfur, q is 2 and (R^c)_q for a compound corresponds in each case to a row of Table C.

15

Table 359

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ is hydrogen and R² is tert-butyl, E' is sulfur, q is 2 and (R^c)_q for a compound corresponds in each case to a row of Table C.

20

Table 360

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ is hydrogen and R² is phenyl, E' is sulfur, q is 2 and (R^c)_q for a compound corresponds in each case to a row of Table C.

25

Table 361

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ and R² are hydrogen, E' is NH and (R^c)_q for a compound corresponds in each case to a row of Table C.

30

Table 362

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is methyl, E' is NH and (R^c)_q for a compound corresponds in each case to a row of Table C.

35

Table 363

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is ethyl, E' is NH and (R^c)_q for a compound corresponds in each case to a row of Table C.

40

Table 364

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is i-propyl, E' is NH and (R^c)_q for a compound corresponds in each case to a row of Table C.

45

Table 365

Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is

70

hydrogen and R² is n-propyl, E' is NH and (R^c)_q for a compound corresponds in each case to a row of Table C.

Table 366

- 5 Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is cyclo-propyl, E' is NH and (R^c)_q for a compound corresponds in each case to a row of Table C.

Table 367

- 10 Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is n-butyl, E' is NH and (R^c)_q for a compound corresponds in each case to a row of Table C.

Table 368

- 15 Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is i-butyl, E' is NH and (R^c)_q for a compound corresponds in each case to a row of Table C.

Table 369

- 20 Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is tert-butyl, E' is NH and (R^c)_q for a compound corresponds in each case to a row of Table C.

Table 370

- 25 Compounds of the formula I.6 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is phenyl, E' is NH and (R^c)_q for a compound corresponds in each case to a row of Table C.

Table 371

- 30 Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ and R² are hydrogen, E' is NH and (R^c)_q for a compound corresponds in each case to a row of Table C.

Table 372

- 35 Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is hydrogen and R² is methyl, E' is NH and (R^c)_q for a compound corresponds in each case to a row of Table C.

Table 373

- 40 Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is hydrogen and R² is ethyl, E' is NH and (R^c)_q for a compound corresponds in each case to a row of Table C.

Table 374

- 45 Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is hydrogen and R² is i-propyl, E' is NH and (R^c)_q for a compound

71

corresponds in each case to a row of Table C.

Table 375

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is 5 hydrogen and R² is n-propyl, E' is NH and (R^c)_q for a compound corresponds in each case to a row of Table C.

Table 376

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is 10 hydrogen and R² is cyclo-propyl, E' is NH and (R^c)_q for a compound corresponds in each case to a row of Table C.

Table 377

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is 15 hydrogen and R² is n-butyl, E' is NH and (R^c)_q for a compound corresponds in each case to a row of Table C.

Table 378

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is 20 hydrogen and R² is i-butyl, E' is NH and (R^c)_q for a compound corresponds in each case to a row of Table C.

Table 379

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is 25 hydrogen and R² is tert-butyl, E' is NH and (R^c)_q for a compound corresponds in each case to a row of Table C.

Table 380

Compounds of the formula I.6 wherein X is hydrogen, p is 4, R¹ is 30 hydrogen and R² is phenyl, E' is NH and (R^c)_q for a compound corresponds in each case to a row of Table C.

Table 381

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ and 35 R² are hydrogen, E' is NH and (R^c)_q for a compound corresponds in each case to a row of Table C.

Table 382

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is 40 hydrogen and R² is methyl, E' is NH and (R^c)_q for a compound corresponds in each case to a row of Table C.

Table 383

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is 45 hydrogen and R² is ethyl, E' is NH and (R^c)_q for a compound corresponds in each case to a row of Table C.

Table 384

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is hydrogen and R² is i-propyl, E' is NH and (R^c)_q for a compound corresponds in each case to a row of Table C.

5

Table 385

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is hydrogen and R² is n-propyl, E' is NH and (R^c)_q for a compound corresponds in each case to a row of Table C.

10

Table 386

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is hydrogen and R² is cyclo-propyl, E' is NH and (R^c)_q for a compound corresponds in each case to a row of Table C.

15

Table 387

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is hydrogen and R² is n-butyl, E' is NH and (R^c)_q for a compound corresponds in each case to a row of Table C.

20

Table 388

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is hydrogen and R² is i-butyl, E' is NH and (R^c)_q for a compound corresponds in each case to a row of Table C.

25

Table 389

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is hydrogen and R² is tert-butyl, E' is NH and (R^c)_q for a compound corresponds in each case to a row of Table C.

30

Table 390

Compounds of the formula I.6 wherein X is fluorine, p is 2, R¹ is hydrogen and R² is phenyl, E' is NH and (R^c)_q for a compound corresponds in each case to a row of Table C.

35

Table 391

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ and R² are hydrogen, E' is NH and (R^c)_q for a compound corresponds in each case to a row of Table C.

40

Table 392

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ is hydrogen and R² is methyl, E' is NH and (R^c)_q for a compound corresponds in each case to a row of Table C.

45

Table 393

Compounds of the formula I.6 wherein X is fluorine, p is 4, R¹ is

73

hydrogen and R^2 is ethyl, E' is NH and $(R^c)_q$ for a compound corresponds in each case to a row of Table C.

Table 394

- 5 Compounds of the formula I.6 wherein X is fluorine, p is 4, R^1 is hydrogen and R^2 is i-propyl, E' is NH and $(R^c)_q$ for a compound corresponds in each case to a row of Table C.

Table 395

- 10 Compounds of the formula I.6 wherein X is fluorine, p is 4, R^1 is hydrogen and R^2 is n-propyl, E' is NH and $(R^c)_q$ for a compound corresponds in each case to a row of Table C.

Table 396

- 15 Compounds of the formula I.6 wherein X is fluorine, p is 4, R^1 is hydrogen and R^2 is cyclo-propyl, E' is NH and $(R^c)_q$ for a compound corresponds in each case to a row of Table C.

Table 397

- 20 Compounds of the formula I.6 wherein X is fluorine, p is 4, R^1 is hydrogen and R^2 is n-butyl, E' is NH and $(R^c)_q$ for a compound corresponds in each case to a row of Table C.

Table 398

- 25 Compounds of the formula I.6 wherein X is fluorine, p is 4, R^1 is hydrogen and R^2 is i-butyl, E' is NH and $(R^c)_q$ for a compound corresponds in each case to a row of Table C.

Table 399

- 30 Compounds of the formula I.6 wherein X is fluorine, p is 4, R^1 is hydrogen and R^2 is tert-butyl, E' is NH and $(R^c)_q$ for a compound corresponds in each case to a row of Table C.

Table 400

- 35 Compounds of the formula I.6 wherein X is fluorine, p is 4, R^1 is hydrogen and R^2 is phenyl, E' is NH and $(R^c)_q$ for a compound corresponds in each case to a row of Table C.




Table C

Nr.	R^c (4-Position)	R^c (5-Position)
45 C-1	CH ₃	CH ₃
C-2	CH ₂ CH ₃	CH ₃
C-3	CH ₂ CH ₂ CH ₃	CH ₃

	Nr.	R ^c (4-Position)	R ^c (5-Position)
	C-4	CH(CH ₃) ₂	CH ₃
	C-5	C(CH ₃) ₃	CH ₃
5	C-6	Cl	CH ₃
	C-7	Br	CH ₃
	C-8	CH ₂ OCH ₃	CH ₃
	C-9	COOCH ₃	CH ₃
10	C-10	C ₆ H ₅	CH ₃
	C-11	2-F-C ₆ H ₄	CH ₃
	C-12	3-F-C ₆ H ₄	CH ₃
	C-13	4-F-C ₆ H ₄	CH ₃
	C-14	2-Cl-C ₆ H ₄	CH ₃
15	C-15	3-Cl-C ₆ H ₄	CH ₃
	C-16	4-Cl-C ₆ H ₄	CH ₃
	C-17	2-CH ₃ -C ₆ H ₄	CH ₃
	C-18	3-CH ₃ -C ₆ H ₄	CH ₃
20	C-19	4-CH ₃ -C ₆ H ₄	CH ₃
	C-20	CH ₃	CH ₂ CH ₃
	C-21	CH ₂ CH ₃	CH ₂ CH ₃
	C-22	CH ₂ CH ₂ CH ₃	CH ₂ CH ₃
25	C-23	CH(CH ₃) ₂	CH ₂ CH ₃
	C-24	C(CH ₃) ₃	CH ₂ CH ₃
	C-25	Cl	CH ₂ CH ₃
	C-26	Br	CH ₂ CH ₃
	C-27	CH ₂ OCH ₃	CH ₂ CH ₃
30	C-28	COOCH ₃	CH ₂ CH ₃
	C-29	C ₆ H ₅	CH ₂ CH ₃
	C-30	2-F-C ₆ H ₄	CH ₂ CH ₃
	C-31	3-F-C ₆ H ₄	CH ₂ CH ₃
35	C-32	4-F-C ₆ H ₄	CH ₂ CH ₃
	C-33	2-Cl-C ₆ H ₄	CH ₂ CH ₃
	C-34	3-Cl-C ₆ H ₄	CH ₂ CH ₃
	C-35	4-Cl-C ₆ H ₄	CH ₂ CH ₃
40	C-36	2-CH ₃ -C ₆ H ₄	CH ₂ CH ₃
	C-37	3-CH ₃ -C ₆ H ₄	CH ₂ CH ₃
	C-38	4-CH ₃ -C ₆ H ₄	CH ₂ CH ₃
	C-39	CH ₃	CH ₂ CH ₂ CH ₃
45	C-40	CH ₂ CH ₃	CH ₂ CH ₂ CH ₃
	C-41	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ CH ₃
	C-42	CH(CH ₃) ₂	CH ₂ CH ₂ CH ₃

	Nr.	R ^c (4-Position)	R ^c (5-Position)
5	C-43	C(CH ₃) ₃	CH ₂ CH ₂ CH ₃
	C-44	Cl	CH ₂ CH ₂ CH ₃
	C-45	Br	CH ₂ CH ₂ CH ₃
	C-46	CH ₂ OCH ₃	CH ₂ CH ₂ CH ₃
	C-47	COOCH ₃	CH ₂ CH ₂ CH ₃
10	C-48	C ₆ H ₅	CH ₂ CH ₂ CH ₃
	C-49	2-F-C ₆ H ₄	CH ₂ CH ₂ CH ₃
	C-50	3-F-C ₆ H ₄	CH ₂ CH ₂ CH ₃
	C-51	4-F-C ₆ H ₄	CH ₂ CH ₂ CH ₃
	C-52	2-Cl-C ₆ H ₄	CH ₂ CH ₂ CH ₃
15	C-53	3-Cl-C ₆ H ₄	CH ₂ CH ₂ CH ₃
	C-54	4-Cl-C ₆ H ₄	CH ₂ CH ₂ CH ₃
	C-55	2-CH ₃ -C ₆ H ₄	CH ₂ CH ₂ CH ₃
	C-56	3-CH ₃ -C ₆ H ₄	CH ₂ CH ₂ CH ₃
	C-57	4-CH ₃ -C ₆ H ₄	CH ₂ CH ₂ CH ₃
20	C-58	CH ₃	CH(CH ₃) ₂
	C-59	CH ₂ CH ₃	CH(CH ₃) ₂
	C-60	CH ₂ CH ₂ CH ₃	CH(CH ₃) ₂
	C-61	CH(CH ₃) ₂	CH(CH ₃) ₂
	C-62	C(CH ₃) ₃	CH(CH ₃) ₂
25	C-63	Cl	CH(CH ₃) ₂
	C-64	Br	CH(CH ₃) ₂
	C-65	CH ₂ OCH ₃	CH(CH ₃) ₂
	C-66	COOCH ₃	CH(CH ₃) ₂
	C-67	C ₆ H ₅	CH(CH ₃) ₂
30	C-68	2-F-C ₆ H ₄	CH(CH ₃) ₂
	C-69	3-F-C ₆ H ₄	CH(CH ₃) ₂
	C-70	4-F-C ₆ H ₄	CH(CH ₃) ₂
	C-71	2-Cl-C ₆ H ₄	CH(CH ₃) ₂
	C-72	3-Cl-C ₆ H ₄	CH(CH ₃) ₂
35	C-73	4-Cl-C ₆ H ₄	CH(CH ₃) ₂
	C-74	2-CH ₃ -C ₆ H ₄	CH(CH ₃) ₂
	C-75	3-CH ₃ -C ₆ H ₄	CH(CH ₃) ₂
	C-76	4-CH ₃ -C ₆ H ₄	CH(CH ₃) ₂
	C-77	CH ₃	C(CH ₃) ₃
40	C-78	CH ₂ CH ₃	C(CH ₃) ₃
	C-79	CH ₂ CH ₂ CH ₃	C(CH ₃) ₃
	C-80	CH(CH ₃) ₂	C(CH ₃) ₃
	C-81	C(CH ₃) ₃	C(CH ₃) ₃

	Nr.	R ^c (4-Position)	R ^c (5-Position)
	C-82	Cl	C(CH ₃) ₃
	C-83	Br	C(CH ₃) ₃
5	C-84	CH ₂ OCH ₃	C(CH ₃) ₃
	C-85	COOCH ₃	C(CH ₃) ₃
	C-86	C ₆ H ₅	C(CH ₃) ₃
	C-87	2-F-C ₆ H ₄	C(CH ₃) ₃
10	C-88	3-F-C ₆ H ₄	C(CH ₃) ₃
	C-89	4-F-C ₆ H ₄	C(CH ₃) ₃
	C-90	2-Cl-C ₆ H ₄	C(CH ₃) ₃
	C-91	3-Cl-C ₆ H ₄	C(CH ₃) ₃
	C-92	4-Cl-C ₆ H ₄	C(CH ₃) ₃
15	C-93	2-CH ₃ -C ₆ H ₄	C(CH ₃) ₃
	C-94	3-CH ₃ -C ₆ H ₄	C(CH ₃) ₃
	C-95	4-CH ₃ -C ₆ H ₄	C(CH ₃) ₃
	C-96	CH ₃	Cl
20	C-97	CH ₂ CH ₃	Cl
	C-98	CH ₂ CH ₂ CH ₃	Cl
	C-99	CH(CH ₃) ₂	Cl
	C-100	C(CH ₃) ₃	Cl
25	C-101	Cl	Cl
	C-102	Br	Cl
	C-103	CH ₂ OCH ₃	Cl
	C-104	COOCH ₃	Cl
	C-105	C ₆ H ₅	Cl
30	C-106	2-F-C ₆ H ₄	Cl
	C-107	3-F-C ₆ H ₄	Cl
	C-108	4-F-C ₆ H ₄	Cl
	C-109	2-Cl-C ₆ H ₄	Cl
35	C-110	3-Cl-C ₆ H ₄	Cl
	C-111	4-Cl-C ₆ H ₄	Cl
	C-112	2-CH ₃ -C ₆ H ₄	Cl
	C-113	3-CH ₃ -C ₆ H ₄	Cl
40	C-114	4-CH ₃ -C ₆ H ₄	Cl
	C-115	CH ₃	Br
	C-116	CH ₂ CH ₃	Br
	C-117	CH ₂ CH ₂ CH ₃	Br
45	C-118	CH(CH ₃) ₂	Br
	C-119	C(CH ₃) ₃	Br
	C-120	Cl	Br

	Nr.	R ^c (4-Position)	R ^c (5-Position)
5	C-121	Br	Br
	C-122	CH ₂ OCH ₃	Br
	C-123	COOCH ₃	Br
	C-124	C ₆ H ₅	Br
	C-125	2-F-C ₆ H ₄	Br
10	C-126	3-F-C ₆ H ₄	Br
	C-127	4-F-C ₆ H ₄	Br
	C-128	2-Cl-C ₆ H ₄	Br
	C-129	3-Cl-C ₆ H ₄	Br
	C-130	4-Cl-C ₆ H ₄	Br
15	C-131	2-CH ₃ -C ₆ H ₄	Br
	C-132	3-CH ₃ -C ₆ H ₄	Br
	C-133	4-CH ₃ -C ₆ H ₄	Br
	C-134	CH ₃	CH ₂ OCH ₃
	C-135	CH ₂ CH ₃	CH ₂ OCH ₃
20	C-136	CH ₂ CH ₂ CH ₃	CH ₂ OCH ₃
	C-137	CH(CH ₃) ₂	CH ₂ OCH ₃
	C-138	C(CH ₃) ₃	CH ₂ OCH ₃
	C-139	Cl	CH ₂ OCH ₃
	C-140	Br	CH ₂ OCH ₃
25	C-141	CH ₂ OCH ₃	CH ₂ OCH ₃
	C-142	COOCH ₃	CH ₂ OCH ₃
	C-143	C ₆ H ₅	CH ₂ OCH ₃
	C-144	2-F-C ₆ H ₄	CH ₂ OCH ₃
	C-145	3-F-C ₆ H ₄	CH ₂ OCH ₃
30	C-146	4-F-C ₆ H ₄	CH ₂ OCH ₃
	C-147	2-Cl-C ₆ H ₄	CH ₂ OCH ₃
	C-148	3-Cl-C ₆ H ₄	CH ₂ OCH ₃
	C-149	4-Cl-C ₆ H ₄	CH ₂ OCH ₃
	C-150	2-CH ₃ -C ₆ H ₄	CH ₂ OCH ₃
35	C-151	3-CH ₃ -C ₆ H ₄	CH ₂ OCH ₃
	C-152	4-CH ₃ -C ₆ H ₄	CH ₂ OCH ₃
	C-153	CH ₃	COOCH ₃
	C-154	CH ₂ CH ₃	COOCH ₃
	C-155	CH ₂ CH ₂ CH ₃	COOCH ₃
40	C-156	CH(CH ₃) ₂	COOCH ₃
	C-157	C(CH ₃) ₃	COOCH ₃
	C-158	Cl	COOCH ₃
	C-159	Br	COOCH ₃
	C-159	Br	COOCH ₃

	Nr.	R ^c (4-Position)	R ^c (5-Position)
	C-160	CH ₂ OCH ₃	COOCH ₃
	C-161	COOCH ₃	COOCH ₃
5	C-162	C ₆ H ₅	COOCH ₃
	C-163	2-F-C ₆ H ₄	COOCH ₃
	C-164	3-F-C ₆ H ₄	COOCH ₃
	C-165	4-F-C ₆ H ₄	COOCH ₃
10	C-166	2-Cl-C ₆ H ₄	COOCH ₃
	C-167	3-Cl-C ₆ H ₄	COOCH ₃
	C-168	4-Cl-C ₆ H ₄	COOCH ₃
	C-169	2-CH ₃ -C ₆ H ₄	COOCH ₃
	C-170	3-CH ₃ -C ₆ H ₄	COOCH ₃
15	C-171	4-CH ₃ -C ₆ H ₄	COOCH ₃
	C-172	CH ₃	C ₆ H ₅
	C-173	CH ₂ CH ₃	C ₆ H ₅
	C-174	CH ₂ CH ₂ CH ₃	C ₆ H ₅
20	C-175	CH(CH ₃) ₂	C ₆ H ₅
	C-176	C(CH ₃) ₃	C ₆ H ₅
	C-177	Cl	C ₆ H ₅
	C-178	Br	C ₆ H ₅
25	C-179	CH ₂ OCH ₃	C ₆ H ₅
	C-180	COOCH ₃	C ₆ H ₅
	C-181	C ₆ H ₅	C ₆ H ₅
	C-182	2-F-C ₆ H ₄	C ₆ H ₅
30	C-183	3-F-C ₆ H ₄	C ₆ H ₅
	C-184	4-F-C ₆ H ₄	C ₆ H ₅
	C-185	2-Cl-C ₆ H ₄	C ₆ H ₅
	C-186	3-Cl-C ₆ H ₄	C ₆ H ₅
	C-187	4-Cl-C ₆ H ₄	C ₆ H ₅
35	C-188	2-CH ₃ -C ₆ H ₄	C ₆ H ₅
	C-189	3-CH ₃ -C ₆ H ₄	C ₆ H ₅
	C-190	4-CH ₃ -C ₆ H ₄	C ₆ H ₅
	C-191	CH ₃	2-F-C ₆ H ₄
40	C-192	CH ₂ CH ₃	2-F-C ₆ H ₄
	C-193	CH ₂ CH ₂ CH ₃	2-F-C ₆ H ₄
	C-194	CH(CH ₃) ₂	2-F-C ₆ H ₄
	C-195	C(CH ₃) ₃	2-F-C ₆ H ₄
45	C-196	Cl	2-F-C ₆ H ₄
	C-197	Br	2-F-C ₆ H ₄
	C-198	CH ₂ OCH ₃	2-F-C ₆ H ₄

	Nr.	R ^c (4-Position)	R ^c (5-Position)
	C-199	COOCH ₃	2-F-C ₆ H ₄
	C-200	C ₆ H ₅	2-F-C ₆ H ₄
5	C-201	2-F-C ₆ H ₄	2-F-C ₆ H ₄
	C-202	3-F-C ₆ H ₄	2-F-C ₆ H ₄
	C-203	4-F-C ₆ H ₄	2-F-C ₆ H ₄
	C-204	2-Cl-C ₆ H ₄	2-F-C ₆ H ₄
10	C-205	3-Cl-C ₆ H ₄	2-F-C ₆ H ₄
	C-206	4-Cl-C ₆ H ₄	2-F-C ₆ H ₄
	C-207	2-CH ₃ -C ₆ H ₄	2-F-C ₆ H ₄
	C-208	3-CH ₃ -C ₆ H ₄	2-F-C ₆ H ₄
	C-209	4-CH ₃ -C ₆ H ₄	2-F-C ₆ H ₄
15	C-210	CH ₃	3-F-C ₆ H ₄
	C-211	CH ₂ CH ₃	3-F-C ₆ H ₄
	C-212	CH ₂ CH ₂ CH ₃	3-F-C ₆ H ₄
	C-213	CH(CH ₃) ₂	3-F-C ₆ H ₄
20	C-214	C(CH ₃) ₃	3-F-C ₆ H ₄
	C-215	Cl	3-F-C ₆ H ₄
	C-216	Br	3-F-C ₆ H ₄
	C-217	CH ₂ OCH ₃	3-F-C ₆ H ₄
25	C-218	COOCH ₃	3-F-C ₆ H ₄
	C-219	C ₆ H ₅	3-F-C ₆ H ₄
	C-220	2-F-C ₆ H ₄	3-F-C ₆ H ₄
	C-221	3-F-C ₆ H ₄	3-F-C ₆ H ₄
	C-222	4-F-C ₆ H ₄	3-F-C ₆ H ₄
30	C-223	2-Cl-C ₆ H ₄	3-F-C ₆ H ₄
	C-224	3-Cl-C ₆ H ₄	3-F-C ₆ H ₄
	C-225	4-Cl-C ₆ H ₄	3-F-C ₆ H ₄
	C-226	2-CH ₃ -C ₆ H ₄	3-F-C ₆ H ₄
35	C-227	3-CH ₃ -C ₆ H ₄	3-F-C ₆ H ₄
	C-228	4-CH ₃ -C ₆ H ₄	3-F-C ₆ H ₄
	C-229	CH ₃	4-F-C ₆ H ₄
	C-230	CH ₂ CH ₃	4-F-C ₆ H ₄
40	C-231	CH ₂ CH ₂ CH ₃	4-F-C ₆ H ₄
	C-232	CH(CH ₃) ₂	4-F-C ₆ H ₄
	C-233	C(CH ₃) ₃	4-F-C ₆ H ₄
	C-234	Cl	4-F-C ₆ H ₄
	C-235	Br	4-F-C ₆ H ₄
45	C-236	CH ₂ OCH ₃	4-F-C ₆ H ₄
	C-237	COOCH ₃	4-F-C ₆ H ₄

	Nr.	R ^c (4-Position)	R ^c (5-Position)
	C-238	C ₆ H ₅	4-F-C ₆ H ₄
	C-239	2-F-C ₆ H ₄	4-F-C ₆ H ₄
5	C-240	3-F-C ₆ H ₄	4-F-C ₆ H ₄
	C-241	4-F-C ₆ H ₄	4-F-C ₆ H ₄
	C-242	2-Cl-C ₆ H ₄	4-F-C ₆ H ₄
	C-243	3-Cl-C ₆ H ₄	4-F-C ₆ H ₄
10	C-244	4-Cl-C ₆ H ₄	4-F-C ₆ H ₄
	C-245	2-CH ₃ -C ₆ H ₄	4-F-C ₆ H ₄
	C-246	3-CH ₃ -C ₆ H ₄	4-F-C ₆ H ₄
	C-247	4-CH ₃ -C ₆ H ₄	4-F-C ₆ H ₄
15	C-248	CH ₃	2-Cl-C ₆ H ₄
	C-249	CH ₂ CH ₃	2-Cl-C ₆ H ₄
	C-250	CH ₂ CH ₂ CH ₃	2-Cl-C ₆ H ₄
	C-251	CH(CH ₃) ₂	2-Cl-C ₆ H ₄
	C-252	C(CH ₃) ₃	2-Cl-C ₆ H ₄
20	C-253	Cl	2-Cl-C ₆ H ₄
	C-254	Br	2-Cl-C ₆ H ₄
	C-255	CH ₂ OCH ₃	2-Cl-C ₆ H ₄
	C-256	COOCH ₃	2-Cl-C ₆ H ₄
25	C-257	C ₆ H ₅	2-Cl-C ₆ H ₄
	C-258	2-F-C ₆ H ₄	2-Cl-C ₆ H ₄
	C-259	3-F-C ₆ H ₄	2-Cl-C ₆ H ₄
	C-260	4-F-C ₆ H ₄	2-Cl-C ₆ H ₄
30	C-261	2-Cl-C ₆ H ₄	2-Cl-C ₆ H ₄
	C-262	3-Cl-C ₆ H ₄	2-Cl-C ₆ H ₄
	C-263	4-Cl-C ₆ H ₄	2-Cl-C ₆ H ₄
	C-264	2-CH ₃ -C ₆ H ₄	2-Cl-C ₆ H ₄
	C-265	3-CH ₃ -C ₆ H ₄	2-Cl-C ₆ H ₄
35	C-266	4-CH ₃ -C ₆ H ₄	2-Cl-C ₆ H ₄
	C-267	CH ₃	3-Cl-C ₆ H ₄
	C-268	CH ₂ CH ₃	3-Cl-C ₆ H ₄
	C-269	CH ₂ CH ₂ CH ₃	3-Cl-C ₆ H ₄
40	C-270	CH(CH ₃) ₂	3-Cl-C ₆ H ₄
	C-271	C(CH ₃) ₃	3-Cl-C ₆ H ₄
	C-272	Cl	3-Cl-C ₆ H ₄
	C-273	Br	3-Cl-C ₆ H ₄
45	C-274	CH ₂ OCH ₃	3-Cl-C ₆ H ₄
	C-275	COOCH ₃	3-Cl-C ₆ H ₄
	C-276	C ₆ H ₅	3-Cl-C ₆ H ₄

	Nr.	R ^c (4-Position)	R ^c (5-Position)
5	C-277	2-F-C ₆ H ₄	3-Cl-C ₆ H ₄
	C-278	3-F-C ₆ H ₄	3-Cl-C ₆ H ₄
	C-279	4-F-C ₆ H ₄	3-Cl-C ₆ H ₄
	C-280	2-Cl-C ₆ H ₄	3-Cl-C ₆ H ₄
	C-281	3-Cl-C ₆ H ₄	3-Cl-C ₆ H ₄
10	C-282	4-Cl-C ₆ H ₄	3-Cl-C ₆ H ₄
	C-283	2-CH ₃ -C ₆ H ₄	3-Cl-C ₆ H ₄
	C-284	3-CH ₃ -C ₆ H ₄	3-Cl-C ₆ H ₄
	C-285	4-CH ₃ -C ₆ H ₄	3-Cl-C ₆ H ₄
	C-286	CH ₃	4-Cl-C ₆ H ₄
15	C-287	CH ₂ CH ₃	4-Cl-C ₆ H ₄
	C-288	CH ₂ CH ₂ CH ₃	4-Cl-C ₆ H ₄
	C-289	CH(CH ₃) ₂	4-Cl-C ₆ H ₄
	C-290	C(CH ₃) ₃	4-Cl-C ₆ H ₄
	C-291	Cl	4-Cl-C ₆ H ₄
20	C-292	Br	4-Cl-C ₆ H ₄
	C-293	CH ₂ OCH ₃	4-Cl-C ₆ H ₄
	C-294	COOCH ₃	4-Cl-C ₆ H ₄
	C-295	C ₆ H ₅	4-Cl-C ₆ H ₄
	C-296	2-F-C ₆ H ₄	4-Cl-C ₆ H ₄
25	C-297	3-F-C ₆ H ₄	4-Cl-C ₆ H ₄
	C-298	4-F-C ₆ H ₄	4-Cl-C ₆ H ₄
	C-299	2-Cl-C ₆ H ₄	4-Cl-C ₆ H ₄
	C-300	3-Cl-C ₆ H ₄	4-Cl-C ₆ H ₄
	C-301	4-Cl-C ₆ H ₄	4-Cl-C ₆ H ₄
30	C-302	2-CH ₃ -C ₆ H ₄	4-Cl-C ₆ H ₄
	C-303	3-CH ₃ -C ₆ H ₄	4-Cl-C ₆ H ₄
	C-304	4-CH ₃ -C ₆ H ₄	4-Cl-C ₆ H ₄
	C-305	CH ₃	2-CH ₃ -C ₆ H ₄
	C-306	CH ₂ CH ₃	2-CH ₃ -C ₆ H ₄
35	C-307	CH ₂ CH ₂ CH ₃	2-CH ₃ -C ₆ H ₄
	C-308	CH(CH ₃) ₂	2-CH ₃ -C ₆ H ₄
	C-309	C(CH ₃) ₃	2-CH ₃ -C ₆ H ₄
	C-310	Cl	2-CH ₃ -C ₆ H ₄
	C-311	Br	2-CH ₃ -C ₆ H ₄
40	C-312	CH ₂ OCH ₃	2-CH ₃ -C ₆ H ₄
	C-313	COOCH ₃	2-CH ₃ -C ₆ H ₄
	C-314	C ₆ H ₅	2-CH ₃ -C ₆ H ₄
	C-315	2-F-C ₆ H ₄	2-CH ₃ -C ₆ H ₄
	C-315	2-F-C ₆ H ₄	2-CH ₃ -C ₆ H ₄

	Nr.	R ^c (4-Position)	R ^c (5-Position)
	C-316	3-F-C ₆ H ₄	2-CH ₃ -C ₆ H ₄
	C-317	4-F-C ₆ H ₄	2-CH ₃ -C ₆ H ₄
5	C-318	2-Cl-C ₆ H ₄	2-CH ₃ -C ₆ H ₄
	C-319	3-Cl-C ₆ H ₄	2-CH ₃ -C ₆ H ₄
	C-320	4-Cl-C ₆ H ₄	2-CH ₃ -C ₆ H ₄
	C-321	2-CH ₃ -C ₆ H ₄	2-CH ₃ -C ₆ H ₄
10	C-322	3-CH ₃ -C ₆ H ₄	2-CH ₃ -C ₆ H ₄
	C-323	4-CH ₃ -C ₆ H ₄	2-CH ₃ -C ₆ H ₄
	C-324	CH ₃	3-CH ₃ -C ₆ H ₄
	C-325	CH ₂ CH ₃	3-CH ₃ -C ₆ H ₄
	C-326	CH ₂ CH ₂ CH ₃	3-CH ₃ -C ₆ H ₄
15	C-327	CH(CH ₃) ₂	3-CH ₃ -C ₆ H ₄
	C-328	C(CH ₃) ₃	3-CH ₃ -C ₆ H ₄
	C-329	Cl	3-CH ₃ -C ₆ H ₄
	C-330	Br	3-CH ₃ -C ₆ H ₄
20	C-331	CH ₂ OCH ₃	3-CH ₃ -C ₆ H ₄
	C-332	COOCH ₃	3-CH ₃ -C ₆ H ₄
	C-333	C ₆ H ₅	3-CH ₃ -C ₆ H ₄
	C-334	2-F-C ₆ H ₄	3-CH ₃ -C ₆ H ₄
25	C-335	3-F-C ₆ H ₄	3-CH ₃ -C ₆ H ₄
	C-336	4-F-C ₆ H ₄	3-CH ₃ -C ₆ H ₄
	C-337	2-Cl-C ₆ H ₄	3-CH ₃ -C ₆ H ₄
	C-338	3-Cl-C ₆ H ₄	3-CH ₃ -C ₆ H ₄
	C-339	4-Cl-C ₆ H ₄	3-CH ₃ -C ₆ H ₄
30	C-340	2-CH ₃ -C ₆ H ₄	3-CH ₃ -C ₆ H ₄
	C-341	3-CH ₃ -C ₆ H ₄	3-CH ₃ -C ₆ H ₄
	C-342	4-CH ₃ -C ₆ H ₄	3-CH ₃ -C ₆ H ₄
	C-343	CH ₃	4-CH ₃ -C ₆ H ₄
35	C-344	CH ₂ CH ₃	4-CH ₃ -C ₆ H ₄
	C-345	CH ₂ CH ₂ CH ₃	4-CH ₃ -C ₆ H ₄
	C-346	CH(CH ₃) ₂	4-CH ₃ -C ₆ H ₄
	C-347	C(CH ₃) ₃	4-CH ₃ -C ₆ H ₄
40	C-348	Cl	4-CH ₃ -C ₆ H ₄
	C-349	Br	4-CH ₃ -C ₆ H ₄
	C-350	CH ₂ OCH ₃	4-CH ₃ -C ₆ H ₄
	C-351	COOCH ₃	4-CH ₃ -C ₆ H ₄
45	C-352	C ₆ H ₅	4-CH ₃ -C ₆ H ₄
	C-353	2-F-C ₆ H ₄	4-CH ₃ -C ₆ H ₄
	C-354	3-F-C ₆ H ₄	4-CH ₃ -C ₆ H ₄

Nr.	R ^c (4-Position)	R ^c (5-Position)
C-355	4-F-C ₆ H ₄	4-CH ₃ -C ₆ H ₄
C-356	2-Cl-C ₆ H ₄	4-CH ₃ -C ₆ H ₄
C-357	3-Cl-C ₆ H ₄	4-CH ₃ -C ₆ H ₄
C-358	4-Cl-C ₆ H ₄	4-CH ₃ -C ₆ H ₄
C-359	2-CH ₃ -C ₆ H ₄	4-CH ₃ -C ₆ H ₄
C-360	3-CH ₃ -C ₆ H ₄	4-CH ₃ -C ₆ H ₄
C-361	4-CH ₃ -C ₆ H ₄	4-CH ₃ -C ₆ H ₄

Table 401

Compounds of the formula I.7 wherein X is hydrogen, p is 2, R¹ and R² are hydrogen and the combination of E and R^c for a compound corresponds in each case to a row of Table D.




Table 402

Compounds of the formula I.7 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is methyl and the combination of E and R^c for a compound corresponds in each case to a row of Table D.

Table 403

Compounds of the formula I.7 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is ethyl and the combination of E and R^c for a compound corresponds in each case to a row of Table D.

Table 404

Compounds of the formula I.7 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is i-propyl and the combination of E and R^c for a compound corresponds in each case to a row of Table D.

Table 405

Compounds of the formula I.7 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is n-propyl and the combination of E and R^c for a compound corresponds in each case to a row of Table D.

Table 406

Compounds of the formula I.7 wherein X is hydrogen, p is 2, R¹ is hydrogen and R² is cyclo-propyl and the combination of E and R^c for a compound corresponds in each case to a row of Table D.

Table 407

Compounds of the formula I.7 wherein X is hydrogen, p is 2, R¹ is

84

hydrogen and R^2 is n-butyl and the combination of E and R^c for a compound corresponds in each case to a row of Table D.

Table 408

- 5 Compounds of the formula I.7 wherein X is hydrogen, p is 2, R^1 is hydrogen and R^2 is i-butyl and the combination of E and R^c for a compound corresponds in each case to a row of Table D.

Table 409

- 10 Compounds of the formula I.7 wherein X is hydrogen, p is 2, R^1 is hydrogen and R^2 is tert-butyl and the combination of E and R^b for a compound corresponds in each case to a row of Table D.

Table 410

- 15 Compounds of the formula I.7 wherein X is hydrogen, p is 2, R^1 is hydrogen and R^2 is phenyl and the combination of E and R^c for a compound corresponds in each case to a row of Table D.

Table 411

- 20 Compounds of the formula I.7 wherein X is hydrogen, p is 4, R^1 and R^2 are hydrogen and the combination of E and R^c for a compound corresponds in each case to a row of Table D.

Table 412

- 25 Compounds of the formula I.7 wherein X is hydrogen, p is 4, R^1 is hydrogen and R^2 is methyl and the combination of E and R^c for a compound corresponds in each case to a row of Table D.

Table 413

- 30 Compounds of the formula I.7 wherein X is hydrogen, p is 4, R^1 is hydrogen and R^2 is ethyl and the combination of E and R^c for a compound corresponds in each case to a row of Table D.

Table 414

- 35 Compounds of the formula I.7 wherein X is hydrogen, p is 4, R^1 is hydrogen and R^2 is i-propyl and the combination of E and R^c for a compound corresponds in each case to a row of Table D.

Table 415

- 40 Compounds of the formula I.7 wherein X is hydrogen, p is 4, R^1 is hydrogen and R^2 is n-propyl and the combination of E and R^c for a compound corresponds in each case to a row of Table D.

Table 416

- 45 Compounds of the formula I.7 wherein X is hydrogen, p is 4, R^1 is hydrogen and R^2 is cyclo-propyl and the combination of E and R^c

85

for a compound corresponds in each case to a row of Table D.

Table 417

Compounds of the formula I.7 wherein X is hydrogen, p is 4, R¹ is
5 hydrogen and R² is n-butyl and the combination of E and R^c for a
compound corresponds in each case to a row of Table D.

Table 418

Compounds of the formula I.7 wherein X is hydrogen, p is 4, R¹ is
10 hydrogen and R² is i-butyl and the combination of E and R^c for a
compound corresponds in each case to a row of Table D.

Table 419

Compounds of the formula I.7 wherein X is hydrogen, p is 4, R¹ is
15 hydrogen and R² is tert-butyl and the combination of E and R^c for
a compound corresponds in each case to a row of Table D.

Table 420

Compounds of the formula I.7 wherein X is hydrogen, p is 4, R¹ is
20 hydrogen and R² is phenyl and the combination of E and R^c for a
compound corresponds in each case to a row of Table D.

Table 421

Compounds of the formula I.7 wherein X is fluorine, p is 2, R¹ and
25 R² are hydrogen and the combination of E and R^c for a compound
corresponds in each case to a row of Table D.

Table 422

Compounds of the formula I.7 wherein X is fluorine, p is 2, R¹ is
30 hydrogen and R² is methyl and the combination of E and R^c for a
compound corresponds in each case to a row of Table D.

Table 423

Compounds of the formula I.7 wherein X is fluorine, p is 2, R¹ is
35 hydrogen and R² is ethyl and the combination of E and R^c for a
compound corresponds in each case to a row of Table D.

Table 424

Compounds of the formula I.7 wherein X is fluorine, p is 2, R¹ is
40 hydrogen and R² is i-propyl and the combination of E and R^c for a
compound corresponds in each case to a row of Table D.

Table 425

Compounds of the formula I.7 wherein X is fluorine, p is 2, R¹ is
45 hydrogen and R² is n-propyl and the combination of E and R^c for a
compound corresponds in each case to a row of Table D.

Table 426

Compounds of the formula I.7 wherein X is fluorine, p is 2, R¹ is hydrogen and R² is cyclo-propyl and the combination of E and R^c for a compound corresponds in each case to a row of Table D.

5

Table 427

Compounds of the formula I.7 wherein X is fluorine, p is 2, R¹ is hydrogen and R² is n-butyl and the combination of E and R^c for a compound corresponds in each case to a row of Table D.

10

Table 428

Compounds of the formula I.7 wherein X is fluorine, p is 2, R¹ is hydrogen and R² is i-butyl and the combination of E and R^c for a compound corresponds in each case to a row of Table D.

15

Table 429

Compounds of the formula I.7 wherein X is fluorine, p is 2, R¹ is hydrogen and R² is tert-butyl and the combination of E and R^c for a compound corresponds in each case to a row of Table D.

20

Table 430

Compounds of the formula I.7 wherein X is fluorine, p is 2, R¹ is hydrogen and R² is phenyl and the combination of E and R^c for a compound corresponds in each case to a row of Table D.

25

Table 431

Compounds of the formula I.7 wherein X is fluorine, p is 4, R¹ and R² are hydrogen and the combination of E and R^c for a compound corresponds in each case to a row of Table D.

30

Table 432

Compounds of the formula I.7 wherein X is fluorine, p is 4, R¹ is hydrogen and R² is methyl and the combination of E and R^c for a compound corresponds in each case to a row of Table D.

35

Table 433

Compounds of the formula I.7 wherein X is fluorine, p is 4, R¹ is hydrogen and R² is ethyl and the combination of E and R^c for a compound corresponds in each case to a row of Table D.

40

Table 434

Compounds of the formula I.7 wherein X is fluorine, p is 4, R¹ is hydrogen and R² is i-propyl and the combination of E and R^c for a compound corresponds in each case to a row of Table D.

45

Table 435

Compounds of the formula I.7 wherein X is fluorine, p is 4, R¹ is

hydrogen and R² is n-propyl and the combination of E and R^c for a compound corresponds in each case to a row of Table D.

Table 436

- 5 Compounds of the formula I.7 wherein X is fluorine, p is 4, R¹ is hydrogen and R² is cyclo-propyl and the combination of E and R^c for a compound corresponds in each case to a row of Table D.

Table 437

- 10 Compounds of the formula I.7 wherein X is fluorine, p is 4, R¹ is hydrogen and R² is n-butyl and the combination of E and R^c for a compound corresponds in each case to a row of Table D.

Table 438

- 15 Compounds of the formula I.7 wherein X is fluorine, p is 4, R¹ is hydrogen and R² is i-butyl and the combination of E and R^c for a compound corresponds in each case to a row of Table D.

Table 439


- 20 Compounds of the formula I.7 wherein X is fluorine, p is 4, R¹ is hydrogen and R² is tert-butyl and the combination of E and R^c for a compound corresponds in each case to a row of Table D.

Table 440

- 25 Compounds of the formula I.7 wherein X is fluorine, p is 4, R¹ is hydrogen and R² is phenyl and the combination of E and R^c for a compound corresponds in each case to a row of Table D.

Table D

30



	Nr.	E	R ^c
35	D-1	O	-
	D-2	O	CH ₃
	D-3	O	CH ₂ CH ₃
	D-4	O	CH ₂ CH ₂ CH ₃
40	D-5	O	CH(CH ₃) ₂
	D-6	O	CH ₂ CH ₂ CH ₂ CH ₃
	D-7	O	CH ₂ CH(CH ₃) ₂
	D-8	O	C(CH ₃) ₃
45	D-9	O	F
	D-10	O	Cl
	D-11	O	Br

	Nr.	E	R ^c
	D-12	O	I
	D-13	O	CF ₃
5	D-14	O	OCF ₃
	D-15	O	OCH ₃
	D-16	O	OCH ₂ CH ₃
	D-17	O	OCH ₂ CH ₂ CH ₃
10	D-18	O	OCH(CH ₃) ₂
	D-19	O	NO ₂
	D-20	O	NH ₂
	D-21	O	NH(C=O)CH ₃
15	D-22	S	-
	D-23	S	CH ₃
	D-24	S	CH ₂ CH ₃
	D-25	S	CH ₂ CH ₂ CH ₃
	D-26	S	CH(CH ₃) ₂
20	D-27	S	CH ₂ CH ₂ CH ₂ CH ₃
	D-28	S	CH ₂ CH(CH ₃) ₂
	D-29	S	C(CH ₃) ₃
	D-30	S	F
25	D-31	S	Cl
	D-32	S	Br
	D-33	S	I
	D-34	S	CF ₃
30	D-35	S	OCF ₃
	D-36	S	OCH ₃
	D-37	S	OCH ₂ CH ₃
	D-38	S	OCH ₂ CH ₂ CH ₃
	D-39	S	OCH(CH ₃) ₂
35	D-40	S	NO ₂
	D-41	S	NH ₂
	D-42	S	NH(C=O)CH ₃

40 The compounds of the formula I are suitable for efficiently controlling nematodes, insects, and arachnids in crop protection. In particular, they are suitable for controlling the following animal pests:

45 insects from the order of the lepidopterans (*Lepidoptera*), for example *Agrotis ypsilon*, *Agrotis segetum*, *Alabama argillacea*, *Anticarsia gemmatalis*, *Argyresthia conjugella*, *Autographa gamma*, *Bupalus piniarius*, *Cacoecia murinana*, *Capua reticulana*, *Cheimato-*

bia brumata, Choristoneura fumiferana, Choristoneura occidenta-
 lis, Cirphis unipuncta, Cydia pomonella, Dendrolimus pini, Dia-
 phania nitidalis, Diatraea grandiosella, Earias insulana, Elasma-
 palpus lignosellus, Eupoecilia ambiguella, Evetria bouliana, Fel-
 5 tia subterranea, Galleria mellonella, Grapholitha funebrana, Gra-
 pholitha molesta, Heliothis armigera, Heliothis virescens, Helio-
 this zea, Hellula undalis, Hibernia defoliaria, Hyphantria cunea,
 Hyponomeuta malinellus, Keiferia lycopersicella, Lambdina fiscel-
 laria, Laphygma exigua, Leucoptera coffeella, Leucoptera sci-
 10 tella, Lithocolletis blancardella, Lobesia botrana, Loxostege
 sticticalis, Lymantria dispar, Lymantria monacha, Lyonetia cler-
 kella, Malacosoma neustria, Mamestra brassicae, Orgyia pseudotsu-
 gata, Ostrinia nubilalis, Panolis flammea, Pectinophora gossy-
 piella, Peridroma saucia, Phalera bucephala, Phthorimaea opercu-
 15 lella, Phyllocnistis citrella, Pieris brassicae, Plathypena sca-
 bra, Plutella xylostella, Pseudoplusia includens, Rhyacionia
 frustrana, Scrobipalpula absoluta, Sitotroga cerealella, Sparga-
 nothis pilleriana, Spodoptera frugiperda, Spodoptera littoralis,
 Spodoptera litura, Thaumatopea pityocampa, Tortrix viridana,
 20 Trichoplusia ni and Zeiraphera canadensis,

beetles (Coleoptera), for example Agrilus sinuatus, Agriotes li-
 neatus, Agriotes obscurus, Amphimallus solstitialis, Anisandrus
 dispar, Anthonomus grandis, Anthonomus pomorum, Atomaria linea-
 25 ris, Blastophagus piniperda, Blitophaga undata, Bruchus rufima-
 nus, Bruchus pisorum, Bruchus lentis, Byctiscus betulae, Cassida
 nebulosa, Cerotoma trifurcata, Ceuthorrhynchus assimilis, Ceu-
 thorrhynchus napi, Chaetocnema tibialis, Conoderus vespertinus,
 Crioceris asparagi, Diabrotica longicornis, Diabrotica 12-puncta-
 30 ta, Diabrotica virgifera, Epilachna varivestis, Epitrix hirtipen-
 nis, Eutinobothrus brasiliensis, Hylobius abietis, Hypera brun-
 neipennis, Hypera postica, Ips typographus, Lema bilineata, Lema
 melanopus, Leptinotarsa decemlineata, Limonius californicus, Lis-
 sorhoptrus oryzophilus, Melanotus communis, Meligethes aeneus,
 35 Melolontha hippocastani, Melolontha melolontha, Oulema oryzae,
 Ortiorrhynchus sulcatus, Otiorrhynchus ovatus, Phaedon cochlea-
 riae, Phyllotreta chrysocephala, Phyllophaga sp., Phyllopertha
 horticola, Phyllotreta nemorum, Phyllotreta striolata, Popillia
 japonica, Sitona lineatus and Sitophilus granaria,

40

dipterans (Diptera), for example Aedes aegypti, Aedes vexans, An-
 astrepha ludens, Anopheles maculipennis, Ceratitis capitata,
 Chrysomya bezziana, Chrysomya hominivorax, Chrysomya macellaria,
 Contarinia sorghicola, Cordylobia anthropophaga, Culex pipiens,
 45 Dacus cucurbitae, Dacus oleae, Dasineura brassicae, Fannia cani-
 cularis, Gasterophilus intestinalis, Glossina morsitans, Haemato-
 bia irritans, Haplodiplosis equestris, Hylemyia platura, Hypoder-

ma lineata, *Liriomyza sativae*, *Liriomyza trifolii*, *Lucilia caprina*, *Lucilia cuprina*, *Lucilia sericata*, *Lycoria pectoralis*, *Mayetiola destructor*, *Musca domestica*, *Muscina stabulans*, *Oestrus ovis*, *Oscinella frit*, *Pegomya hysocyami*, *Phorbia antiqua*, *Phorbia*
 5 *brassicae*, *Phorbia coarctata*, *Rhagoletis cerasi*, *Rhagoletis pomonella*, *Tabanus bovinus*, *Tipula oleracea* and *Tipula paludosa*,

thrips (Thysanoptera), e.g. *Frankliniella fusca*, *Frankliniella occidentalis*, *Frankliniella tritici*, *Scirtothrips citri*, *Thrips*
 10 *oryzae*, *Thrips palmi* and *Thrips tabaci*,

hymenopterans (Hymenoptera), e.g. *Athalia rosae*, *Atta cephalotes*, *Atta sexdens*, *Atta texana*, *Hoplocampa minuta*, *Hoplocampa testudinea*, *Monomorium pharaonis*, *Solenopsis geminata* and *Solenopsis in-*
 15 *victa*,

heteropterans (Heteroptera), e.g. *Acrosternum hilare*, *Blissus leucopterus*, *Cyrtopeltis notatus*, *Dysdercus cingulatus*, *Dysdercus intermedius*, *Eurygaster integriceps*, *Euschistus impictiventris*,
 20 *Leptoglossus phyllopus*, *Lygus lineolaris*, *Lygus pratensis*, *Nezara viridula*, *Piesma quadrata*, *Solubea insularis* and *Thyanta perditor*,

homopterans (Homoptera), e.g. *Acyrtosiphon onobrychis*, *Adelges*
 25 *laricis*, *Aphidula nasturtii*, *Aphis fabae*, *Aphis pomi*, *Aphis sambuci*, *Brachycaudus cardui*, *Brevicoryne brassicae*, *Cerosipha gossypii*, *Dreyfusia nordmannianae*, *Dreyfusia piceae*, *Dysaphis radicola*, *Dysaulacorthum pseudosolani*, *Empoasca fabae*, *Macrosiphum avenae*, *Macrosiphum euphorbiae*, *Macrosiphon rosae*, *Megoura vi-*
 30 *ciae*, *Metopolophium dirhodum*, *Myzodes persicae*, *Myzus cerasi*, *Nilaparvata lugens*, *Pemphigus bursarius*, *Perkinsiella saccharicida*, *Phorodon humuli*, *Psylla mali*, *Psylla piri*, *Rhopalomyzus ascalonicus*, *Rhopalosiphum maidis*, *Sappaphis mala*, *Sappaphis mali*, *Schizaphis graminum*, *Schizoneura lanuginosa*, *Trialeurodes vaporario-*
 35 *rum* and *Viteus vitifolii*,

termites (Isoptera), e.g. *Calotermes flavicollis*, *Leucotermes flavipes*, *Reticulitermes lucifugus* und *Termes natalensis*,

40 orthopterans (Orthoptera), e.g. *Acheta domestica*, *Blatta orientalis*, *Blattella germanica*, *Forficula auricularia*, *Gryllotalpa gryllotalpa*, *Locusta migratoria*, *Melanoplus bivittatus*, *Melanoplus femur-rubrum*, *Melanoplus mexicanus*, *Melanoplus sanguinipes*, *Melanoplus spretus*, *Nomadacris septemfasciata*, *Periplaneta ameri-*
 45 *cana*, *Schistocerca americana*, *Schistocerca peregrina*, *Stauronotus maroccanus* and *Tachycines asynamorus*,

91

Arachnoidea, such as arachnids (Acarina), e.g. *Amblyomma americanum*, *Amblyomma variegatum*, *Argas persicus*, *Boophilus annulatus*, *Boophilus decoloratus*, *Boophilus microplus*, *Brevipalpus phoenicis*, *Bryobia praetiosa*, *Dermacentor silvarum*, *Eotetranychus car-*
 5 *pini*, *Eriophyes sheldoni*, *Hyalomma truncatum*, *Ixodes ricinus*,
Ixodes rubicundus, *Ornithodoros moubata*, *Otobius megnini*, *Parate-*
tetranychus pilosus, *Dermanyssus gallinae*, *Phyllocoptiruta oleivora*,
Polyphagotarsonemus latus, *Psoroptes ovis*, *Rhipicephalus appendi-*
culatus, *Rhipicephalus evertsi*, *Sarcoptes scabiei*, *Tetranychus*
 10 *cinnabarinus*, *Tetranychus kanzawai*, *Tetranychus pacificus*, *Tetra-*
nychus telarius and *Tetranychus urticae*,

Nematodes, especially plant parasitic nematodes such as root knot
 nematodes, *Meloidogyne hapla*, *Meloidogyne incognita*, *Meloidogyne*
 15 *javanica*, and other *Meloidogyne* species; cyst-forming nematodes,
Globodera rostochiensis and other *Globodera* species; *Heterodera*
avenae, *Heterodera glycines*, *Heterodera schachtii*, *Heterodera*
trifolii, and other *Heterodera* species; Seed gall nematodes, *An-*
guina species; Stem and foliar nematodes, *Aphelenchoides* species;
 20 *Sting* nematodes, *Belonolaimus longicaudatus* and other *Belonolai-*
mus species; Pine nematodes, *Bursaphelenchus xylophilus* and other
Bursaphelenchus species; Ring nematodes, *Criconema* species, *Cri-*
conemella species, *Criconemoides* species, *Mesocriconema* species;
 Stem and bulb nematodes, *Ditylenchus destructor*, *Ditylenchus dip-*
 25 *saci* and other *Ditylenchus* species; Awl nematodes, *Dolichodorus*
 species; Spiral nematodes, *Helicotylenchus multicinctus* and
 other *Helicotylenchus* species; Sheath and sheathoid nematodes,
Hemicycliophora species and *Hemicriconemoides* species; *Hirshman-*
niella species; Lance nematodes, *Hoploaimus* species; false root-
 30 knot nematodes, *Nacobbus* species; Needle nematodes, *Longidorus*
elongatus and other *Longidorus* species; Pin nematodes, *Paratylen-*
chus species; Lesion nematodes, *Pratylenchus neglectus*, *Pratylen-*
chus penetrans, *Pratylenchus curvatus*, *Pratylenchus goodeyi*
 and other *Pratylenchus* species; Burrowing nematodes, *Radopholus*
 35 *similis* and other *Radopholus* species; Reniform nematodes, *Roty-*
lenchus robustus and other *Rotylenchus* species; *Scutellonema* spe-
 cies; Stubby root nematodes, *Trichodorus primitivus* and other
Trichodorus species, *Paratrachodorus* species; Stunt nematodes,
Tylenchorhynchus claytoni, *Tylenchorhynchus dubius* and other *Ty-*
 40 *lenchorhynchus* species; Citrus nematodes, *Tylenchulus* species;
 Dagger nematodes, *Xiphinema* species; and other plant parasitic
 nematode species.

The compounds of formula I are especially useful for the control
 45 of nematodes.

92

The compounds I also are suitable for use as fungicides. They exhibit activity against a broad spectrum of phytopathogenic fungi, in particular from the classes of the *Ascomycetes*, *Deuteromycetes*, *Phycomycetes* and *Basidiomycetes*. Some of them act systemically, and they can be employed in crop protection as foliar- and soil-acting fungicides.

They are especially important for controlling a large number of fungi on a variety of crop plants such as wheat, rye, barley, oats, rice, maize, grass, bananas, cotton, soya, coffee, sugar cane, grapevines, fruit species, ornamentals and vegetables such as cucumbers, beans, tomatoes, potatoes and cucurbits, and on the seeds of these plants. Specifically, they are suitable for controlling the following plant diseases:

Alternaria species on vegetables and fruit, *Botrytis cinerea* (gray mold) on strawberries, vegetables, ornamentals and grapevines, *Cercospora arachidicola* on peanuts, *Erysiphe cichoracearum* and *Sphaerotheca fuliginea* on cucurbits, *Erysiphe graminis* (powdery mildew) on cereals, *Fusarium* and *Verticillium* species on various plants, *Helminthosporium* species on cereals, *Mycosphaerella* species on bananas and peanuts, *Phytophthora infestans* on potatoes and tomatoes, *Plasmopara viticola* on grapevines, *Podosphaera leucotricha* on apples, *Pseudocercospora herpotrichoides* on wheat and barley, *Pseudoperonospora* species on hops and cucumbers, *Puccinia* species on cereals, *Pyricularia oryzae* on rice, *Rhizoctonia* species on cotton, rice and lawns, *Septoria nodorum* on wheat, *Uncinula necator* on grapevines, *Ustilago* species on cereals and sugar cane, and *Venturia* species (scab) on apples and pears.

Moreover, the compounds I are suitable for controlling harmful fungi such as *Paecilomyces variotii* in the protection of materials (e.g. wood, paper, paint dispersions, fibers and fabrics) and in the protection of stored products.

The compounds I are applied by treating the fungi, or the plants, seeds, materials or the soil to be protected against fungal attack, with a fungicidally effective amount of the active ingredients. Application can be effected both before and after infection of the materials, plants or seeds by the fungi.

Compounds of formula I are suitable for use as herbicides. Depending upon the application method, compounds I and herbicidal compositions comprising them may be used in crops for the control of unwanted plants. Exemplary are the following crops:

93

Allium cepa, *Ananas comosus*, *Arachis hypogaea*, *Asparagus officinalis*, *Beta vulgaris* spp. *altissima*, *Beta vulgaris* spp. *rapa*, *Brassica napus* var. *napus*, *Brassica napus* var. *napobrassica*, *Brassica rapa* var. *silvestris*, *Camellia sinensis*, *Carthamus tinctorius*, *Carya illinoensis*, *Citrus limon*, *Citrus sinensis*, *Coffea arabica* (*Coffea canephora*, *Coffea liberica*), *Cucumis sativus*, *Cynodon dactylon*, *Daucus carota*, *Elaeis guineensis*, *Fragaria vesca*, *Glycine max*, *Gossypium hirsutum*, (*Gossypium arboreum*, *Gossypium herbaceum*, *Gossypium vitifolium*), *Helianthus annuus*, *Hevea brasiliensis*, *Hordeum vulgare*, *Humulus lupulus*, *Ipomoea batatas*, *Juglans regia*, *Lens culinaris*, *Linum usitatissimum*, *Lycopersicon lycopersicum*, *Malus* spp., *Manihot esculenta*, *Medicago sativa*, *Musa* spp., *Nicotiana tabacum* (*N. rustica*), *Olea europaea*, *Oryza sativa*, *Phaseolus lunatus*, *Phaseolus vulgaris*, *Picea abies*, *Pinus* spp., *Pisum sativum*, *Prunus avium*, *Prunus persica*, *Pyrus communis*, *Ribes sylvestre*, *Ricinus communis*, *Saccharum officinarum*, *Secale cereale*, *Solanum tuberosum*, *Sorghum bicolor* (s. *vulgare*), *Theobroma cacao*, *Trifolium pratense*, *Triticum aestivum*, *Triticum durum*, *Vicia faba*, *Vitis vinifera* und *Zea mays*.

20

Compounds of formula I may also be used in crops that have acquired resistance against other herbicides.

For controlling animal pests, pesticidally active amounts of compounds of formula I are typically applied to the pests or to their food supply, habitat or breeding ground. For the protection of growing plants from attack or infestation by the pests, pesticidally active amounts of the compounds of formula I are typically applied to the foliage, stem or roots of the plants or to the soil or water in which they are growing.

35

The rate of application of active ingredient for controlling animal pests is from 0,01 to 100, preferably 0,1 to 3 kg/ha under field conditions.

The compounds I can be converted into the customary formulations, e.g. solutions, emulsions, suspensions, dusts, powders, pastes and granules. The use form depends on the particular purpose; in any case, it should guarantee a fine and uniform distribution of the compound according to the invention.

45

94

The formulations are prepared in a known manner, e.g. by extending the active ingredient with solvents and/or carriers, if desired using emulsifiers and dispersants, it also being possible to use other organic solvents as auxiliary solvents if water is used as the diluent. Auxiliaries which are suitable are essentially: solvents such as aromatics (e.g. xylene), chlorinated aromatics (e.g. chlorobenzenes), paraffins (e.g. mineral oil fractions), alcohols (e.g. methanol, butanol), ketones (e.g. cyclohexanone), amines (e.g. ethanolamine, dimethylformamide) and water; carriers such as ground natural minerals (e.g. kaolins, clays, talc, chalk) and ground synthetic minerals (e.g. highly-disperse silica, silicates); emulsifiers such as non-ionic and anionic emulsifiers (e.g. polyoxyethylene fatty alcohol ethers, alkylsulfonates and arylsulfonates) and dispersants such as lignin-sulfite waste liquors and methylcellulose.

Suitable surfactants are alkali metal, alkaline earth metal and ammonium salts of lignosulfonic acid, naphthalenesulfonic acid, phenolsulfonic acid, dibutyl-naphthalenesulfonic acid, alkylaryl-sulfonates, alkyl sulfates, alkylsulfonates, fatty alcohol sulfates and fatty acids and their alkali metal and alkaline earth metal salts, salts of sulfated fatty alcohol glycol ether, condensates of sulfonated naphthalene and naphthalene derivatives with formaldehyde, condensates of naphthalene or of naphthalenesulfonic acid with phenol or formaldehyde, polyoxyethylene octylphenyl ether, ethoxylated isooctylphenol, octylphenol, nonylphenol, alkylphenol polyglycol ethers, tributylphenyl polyglycol ethers, alkylaryl polyether alcohols, isotridecyl alcohol, fatty alcohol/ethylene oxide condensates, ethoxylated castor oil, polyoxyethylene alkyl ethers, ethoxylated polyoxypropylene, lauryl alcohol polyglycol ether acetal, sorbitol esters, lignin-sulfite waste liquors and methylcellulose.

Substances which are suitable for the preparation of directly sprayable solutions, emulsions, pastes or oil dispersions are mineral oil fractions of medium to high boiling point, such as kerosene or diesel oil, furthermore coal tar oils and oils of vegetable or animal origin, aliphatic, cyclic and aromatic hydrocarbons, e.g. benzene, toluene, xylene, paraffin, tetrahydronaphthalene, alkylated naphthalenes or their derivatives, methanol, ethanol, propanol, butanol, chloroform, carbon tetrachloride, cyclohexanol, cyclohexanone, chlorobenzene, isophorone, strongly polar solvents, e.g. dimethylformamide, dimethyl sulfoxide, N-methylpyrrolidone and water.

95

Powders, materials for scattering and dusts can be prepared by mixing or concomitantly grinding the active substances with a solid carrier.

5 Granules, e.g. coated granules, impregnated granules and homogeneous granules, can be prepared by binding the active ingredients to solid carriers. Examples of solid carriers are mineral earths, such as silicas, silica gels, silicates, talc, kaolin, attaclay, limestone, lime, chalk, bole, loess, clay, dolomite, diatomaceous
10 earth, calcium sulfate, magnesium sulfate, magnesium oxide, ground synthetic materials, fertilizers, e.g. ammonium sulfate, ammonium phosphate, ammonium nitrate, ureas, and products of vegetable origin, such as cereal meal, tree bark meal, wood meal and nutshell meal, cellulose powders and other solid carriers.

15 In general, the formulations comprise of from 0.01 to 95% by weight, preferably from 0.1 to 90% by weight, of the active ingredient. The active ingredients are employed in a purity of from 90% to 100%, preferably 95% to 100% (according to NMR spectrum).

20 The following are exemplary formulations:

I. 5 parts by weight of a compound according to the invention are mixed intimately with 95 parts by weight of finely divided kaolin. This gives a dust which comprises 5% by weight
25 of the active ingredient.

II. 30 parts by weight of a compound according to the invention are mixed intimately with a mixture of 92 parts by weight of pulverulent silica gel and 8 parts by weight of paraffin oil
30 which had been sprayed onto the surface of this silica gel. This gives a formulation of the active ingredient with good adhesion properties (comprises 23% by weight of active ingredient).

35 III. 10 parts by weight of a compound according to the invention are dissolved in a mixture composed of 90 parts by weight of xylene, 6 parts by weight of the adduct of 8 to 10 mol of ethylene oxide and 1 mol of oleic acid N-monoethanolamide, 2
40 parts by weight of calcium dodecylbenzenesulfonate and 2 parts by weight of the adduct of 40 mol of ethylene oxide and 1 mol of castor oil (comprises 9% by weight of active ingredient).

45

96

- IV. 20 parts by weight of a compound according to the invention are dissolved in a mixture composed of 60 parts by weight of cyclohexanone, 30 parts by weight of isobutanol, 5 parts by weight of the adduct of 7 mol of ethylene oxide and 1 mol of isooctylphenol and 5 parts by weight of the adduct of 40 mol of ethylene oxide and 1 mol of castor oil (comprises 16% by weight of active ingredient).
- 5
- V. 80 parts by weight of a compound according to the invention are mixed thoroughly with 3 parts by weight of sodium diisobutyl-naphthalene- α -sulfonate, 10 parts by weight of the sodium salt of a lignosulfonic acid from a sulfite waste liquor and 7 parts by weight of pulverulent silica gel, and the mixture is ground in a hammer mill (comprises 80% by weight of active ingredient).
- 10
- 15
- VI. 90 parts by weight of a compound according to the invention are mixed with 10 parts by weight of N-methyl- α -pyrrolidone, which gives a solution which is suitable for use in the form of microdrops (comprises 90% by weight of active ingredient).
- 20
- VII. 20 parts by weight of a compound according to the invention are dissolved in a mixture composed of 40 parts by weight of cyclohexanone, 30 parts by weight of isobutanol, 20 parts by weight of the adduct of 7 mol of ethylene oxide and 1 mol of isooctylphenol and 10 parts by weight of the adduct of 40 mol of ethylene oxide and 1 mol of castor oil. Pouring the solution into 100,000 parts by weight of water and finely distributing it therein gives an aqueous dispersion which comprises 0.02% by weight of the active ingredient.
- 25
- 30
- VIII. 20 parts by weight of a compound according to the invention are mixed thoroughly with 3 parts by weight of sodium diisobutyl-naphthalene- α -sulfonate, 17 parts by weight of the sodium salt of a lignosulfonic acid from a sulfite waste liquor and 60 parts by weight of pulverulent silica gel, and the mixture is ground in a hammer mill. Finely distributing the mixture in 20,000 parts by weight of water gives a spray mixture which comprises 0.1% by weight of the active ingredient.
- 35
- 40

The active ingredients can be used as such, in the form of their formulations or the use forms prepared therefrom, e.g. in the form of directly sprayable solutions, powders, suspensions or dispersions, emulsions, oil dispersions, pastes, dusts, materials for spreading, or granules, by means of spraying, atomizing, du-

45

sting, scattering or pouring. The use forms depend entirely on the intended purposes; in any case, this is intended to guarantee the finest possible distribution of the active ingredients according to the invention.

5

Aqueous use forms can be prepared from emulsion concentrates, pastes or wettable powders (sprayable powders, oil dispersions) by adding water. To prepare emulsions, pastes or oil dispersions, the substances as such or dissolved in an oil or solvent, can be
10 homogenized in water by means of wetter, tackifier, dispersant or emulsifier. Alternatively, it is possible to prepare concentrates composed of active substance, wetter, tackifier, dispersant or emulsifier and, if appropriate, solvent or oil, and such concentrates are suitable for dilution with water.

15

The active ingredient concentrations in the ready-to-use products can be varied within substantial ranges. In general, they are from 0.0001 to 10%, preferably from 0.01 to 1%.

20 The active ingredients may also be used successfully in the ultra-low-volume process (ULV), it being possible to apply formulations comprising over 95% by weight of active ingredient, or even the active ingredient without additives.

25 Various types of oils, herbicides, fungicides, other pesticides, or bactericides may be added to the active ingredients, if appropriate also only immediately prior to use (tank mix). These agents can be admixed with the agents according to the invention in a weight ratio of 1:10 to 10:1.

30

In the use form as pesticides in crop protection, the compositions according to the invention can also be present together with other active ingredients, e.g. with herbicides, insecticides, growth regulators, fungicides or else with fertilizers.

35 Mixing the compounds I or the compositions comprising them in the use form as pesticides with other pesticides frequently results in a broader pesticidal spectrum of action.

The following list of pesticides together with which the com-

40 pounds according to the invention can be used, is intended to illustrate the possible combinations.

Organophosphates: Acephate, Azinphos-methyl, Chlorpyrifos, Chlorfenvinphos, Diazinon, Dichlorvos, Dicrotophos, Dimethoate, Disulfoton, Ethion, Fenitrothion, Fenthion, Isoxathion, Malathion, Methamidophos, Methidathion, Methyl-Parathion, Mevinphos, Monocrotophos, Oxydemeton-methyl, Paraoxon, Parathion, Phenthoate, Pho-

98

salone, Phosmet, Phosphamidon, Phorate, Phoxim, Pirimiphos-methyl, Profenofos, Prothiofos, Sulprophos, Triazophos, Trichlorfon;

- 5 Carbamates: Alanycarb, Benfuracarb, Carbaryl, Carbosulfan, Fenoxycarb, Furathiocarb, Indoxacarb, Methiocarb, Methomyl, Oxamyl, Pirimicarb, Propoxur, Thiodicarb, Triazamate;

- Pyrethroids: Bifenthrin, Cyfluthrin, Cypermethrin, Deltamethrin,
10 Esfenvalerate, Ethofenprox, Fenpropathrin, Fenvalerate, Cyhalothrin, Lambda-Cyhalothrin, Permethrin, Silafluofen, Tau-Fluvalinate, Tefluthrin, Tralomethrin, Zeta-Cypermethrin;

- Arthropod growth regulators: a) chitin synthesis inhibitors: benzoylureas: Chlorfluazuron, Diflubenzuron, Flucyclohexuron, Flufenoxuron, Hexaflumuron, Lufenuron, Novaluron, Teflubenzuron, Trifluthuron; Buprofezin, Diofenolan, Hexythiazox, Etoxazole, Clofentazine; b) ecdysone antagonists: Halofenozide, Methoxyfenozide, Tebufenozide; c) juvenoids: Pyriproxyfen, Methoprene, Fenoxycarb;
20 d) lipid biosynthesis inhibitors: Spirodiclofen;

- Various: Abamectin, Acequinocyl, Amitraz, Azadirachtin, Bifenazate, Cartap, Chlorfenapyr, Chlordimeform, Cyromazine, Diafenthiuron, Dinotofuran, Diofenolan, Emamectin, Endosulfan, Fenazaquin, Fipronil, Formetanate, Formetanate, Hydrochloride, Hydramethylnon, Imidacloprid, Indoxacarb, Pyridaben, Pymetrozine, Spinosad, Sulfur, Tebufenpyrad, Thiamethoxam, and Thiocyclam.

- This invention also provides a method for treating, curing, controlling, preventing and protecting warm-blooded animals, including humans, and fish against infestation and infection by helminths, acarids and arthropod endo- and ectoparasites which comprises orally, topically or parenterally administering or applying to said animals an anthelmintically, acaricidally or endo-
30 or ectoparasitically effective amount of compounds of formula I.

- The above method is particularly useful for controlling and preventing helminth, acarid and arthropod endo- and ectoparasitic
40 infestations and infections in warm-blooded animals such as cattle, sheep, swine, camels, deer, horses, poultry, fish, rabbits, goats, mink, fox, chinchillas, rabbits, dogs and cats as well as humans.

- 45 Compounds of formula I are especially useful in controlling helminths and nematodes. Examples for helminths are members of the class Trematoda, commonly known as flukes or flatworms, espe-

cially members of the genera *Fasciola*, *Fascioloides*, *Paramphistomum*, *Dicrocoelium*, *Eurytrema*, *Ophisthorchis*, *Fasciolopsis*, *Echinostoma* and *Paragonimus*. Nematodes which can be controlled by the formula I compounds include the genera *Haemonchus*, *Ostertagia*,
5 *Cooperia*, *Oesphagastomum*, *Nematodirus*, *Dictyocaulus*, *Trichuris*, *Dirofilaria*, *Ancylostoma*, *Ascaris* and the like.

The formula I compounds of this invention also control endoparasitic arthropod infestations such as cattle grub and stomach bot.
10 In addition, acarid and arthropod ectoparasitic infestations in warm-blooded animals and fish including biting lice, sucking lice, bot flies, biting flies, muscoid flies, myiasitic fly larvae, gnats, mosquitoes, fleas, mites, ticks, nasal bots, keds and chiggers may be controlled, prevented or eliminated by the compounds of this invention. Biting lice include members of *Mallophaga* such as *Bovicola bovis*, *Trichodectes canis* and *Damillina ovis*. Sucking lice include members of *Anoplura* such as *Haematopinus eurysternus*, *Haematopinus suis*, *Linognathus vituli* and *Solenopotes capillatus*. Biting flies include members of *Haematobia*. Ticks include *Boophilus*, *Rhipicephalus*, *Ixodes*, *Hyalomma*, *Amblyomma* and *Dermacentor*. The formula I compounds may also be used
20 to control mites which are parasitic on warm-blooded mammals and poultry including mites of the orders *Acariformes* and *Parasitiformes*.

25 For oral administration to warm-blooded animals, the formula I compounds may be formulated as animal feeds, animal feed premixes, animal feed concentrates, pills, solutions, pastes, suspensions, drenches, gels, tablets, boluses and capsules. In addition, the formula I compounds may be administered to the animals
30 in their drinking water. For oral administration, the dosage form chosen should provide the animal with about 0.01 mg/kg to 100 mg/kg of animal body weight per day of the formula I compound.

Alternatively, the formula I compounds may be administered to
35 animals parenterally, for example, by intraruminal, intramuscular, intravenous or subcutaneous injection. The formula I compounds may be dispersed or dissolved in a physiologically acceptable carrier for subcutaneous injection. Alternatively, the formula I compounds may be formulated into an implant for subcutaneous
40 administration. In addition the formula I compound may be transdermally administered to animals. For parenteral administration, the dosage form chosen should provide the animal with about 0.01 mg/kg to 100 mg/kg of animal body weight per day of the formula I compound.

45 The formula I compounds may also be applied topically to the animals in the form of dips, dusts, powders, collars, medallions,

100

sprays and pour-on formulations. For topical application, dips and sprays usually contain about 0.5 ppm to 5,000 ppm and preferably about 1 ppm to 3,000 ppm of the formula I compound. In addition, the formula I compounds may be formulated as ear tags
5 for animals, particularly quadrupeds such as cattle and sheep.

The formula I compounds of this invention may also be used in combination or conjunction with one or more other parasitocidal compounds including anthelmintics, such as benzimidazoles, piperazine, levamisole, pyrantel, and praziquantel; endectocides such
10 as avermectins, and milbemycins; ectoparasitocides such as arylpyrroles, organophosphates, and carbamates, gamabutyric acid inhibitors including fipronil, pyrethroids, spinosads and imidacloprid; insect growth regulators such as pyriproxyfen, and cyromazine; and chitin synthase inhibitors such as benzoylureas including flufenoxuron.
15

The formula I compounds may also be used in combination or conjunction with one or more compounds selected from piperonyl butoxide, N-octyl bicycloheptene dicarboximide, dipropyl pyridine-2,5-dicarboxylate and 1,5a,6,9,9a,9b-hexahydro-4a(4H)-
20 dibenzo-furancarboxaldehyde to broaden the spectrum of activity.

The parasitocidal compositions of the present invention include a
25 parasitocidally effective amount of a formula I compound of this invention or combinations thereof admixed with one or more physiologically tolerable inert, solid or liquid carriers known from veterinary medicinal practice for oral, percutaneous and topical administration. Such compositions may comprise further additives,
30 such as stabilizers, anifoams, viscosity regulators, binders and tackifiers. Whereas commercial products will preferably be formulated as concentrates, the end user will normally employ dilute formulations.

35 Synthesis Examples

With due modification of the starting compounds, the protocols shown in the synthesis examples below were used for obtaining further compounds I. The resulting compounds, together with physical data, are listed in Table I which follows.
40

Example 1

Preparation of 2-(5-Chloro-benzothiazol-2-yl-sulfanyl)-propionic acid ethyl ester

45 A solution of 5-chloro-benzothiazol-2-thiol (1,4 g) in 50 ml acetonitrile was treated with potassium carbonate (1,0 g) and 2-bromo-propionic acid ethyl ester (1,25 g) for 3 hours at room

101

temperature. The solvent was removed in vacuo and the residue redissolved in ethyl acetate and washed with water. The organic extract was dried over anhydrous sodium sulfate. Removal of solvent gave 2,1 g of the title compound as a light-brown oil.

5 $^1\text{H-NMR}$ (δ [CDCl_3]) = 1.3 (m_c , 3H), 1.7 (m_c , 3H), 4.2 (m_c , 2H), 4.7 (m_c , 1H), 7.25 (d, 1H), 7.65 (d, 1H), 7.85 (s, 1H).

Example 2

Preparation of 2-(5-Chloro-benzothiazol-2-yl-sulfanyl)-butyric

10 acid

A solution of 2-(5-chloro-benzothiazol-2-yl-sulfanyl)-butyric acid ethyl ester (1,62 g) in 40 ml of ethanol was treated with 40 ml of water and an aqueous solution of potassium hydroxide in
15 water (85%, 0,36 g) for 14 hours at 25 °C. Ethanol was removed in vacuo, the residue acidified with diluted hydrochloric acid and extracted with ethyl acetate. The organic extract was dried over anhydrous sodium sulfate. Removal of solvent gave 1,5 g of the title compound as a light-brown oil.

20 $^1\text{H-NMR}$ (δ [d_6 -DMSO]) = 1.00 (t, 3H), 2.00 (m_c , 2H), 4.55 (t, 1H), 7.45 (d, 1H), 7.95 (s, 1H), 8.1 (d, 1H).

Example 3

Preparation of 2-[4-(4-chlorophenyl)-thiazol-2-yl-sulfanyl]-

25 hexanoic acid 3,4,4-trifluoro-but-3-enyl ester

A mixture of 2-[4-(4-chlorophenyl)-thiazol-2-yl-sulfanyl]-hexanoic acid (0,3 g), potassium carbonate (0,16 g) and 4-bromo-1,2,2-trifluorobut-1-ene (0,21 g) in 20 ml

30 N,N-dimethylformamide was stirred at 40-45°C for 4 hours. The reaction mixture was poured into ice water and extracted with ethyl acetate. The organic extract was dried over anhydrous sodium sulfate and concentrated in vacuo to give 0,37 g of the title compound as a light-brown oil.

35 $^1\text{H-NMR}$ (δ [CDCl_3]) = 0.9 (t, 3H), 1.15-2.1 (m, 6H), 2.6 (m_c , 2H), 4.25-4.4 (m, 3H), 7.35 (m_c , 3H), 7.8 (d, 2H).

No.	B	D	E	(R ^c) _q	q	R ¹	R ²	m	n	p	Physical data: ¹ H-NMR (δ [ppm])
I.2-16	CH	CH	NH	-	0	H	H	0	0	2	
I.2-17	CH	N	S	-	0	H	H	0	0	2	
I.2-18	N	CH	S	6-CH ₃	1	CH ₂ CH ₃	H	0	0	2	
I.2-19	N	CH	S	6-CH ₃	1	CH(CH ₃) ₂	H	0	0	2	
I.2-20	N	CH	O	-	0	H	H	0	0	2	
I.2-21	N	CH	S	-	0	CH ₃	H	0	0	2	
I.2-22	N	CH	S	6-Br	1	H	H	0	0	2	
I.2-23	CH	CH	S	6-OCH ₂ CH ₃	1	CH ₃	H	0	0	2	
I.2-24	N	CH	S	-	0	H	H	0	0	2	
I.2-25	N	CH	S	-	0	CH(CH ₃) ₂	H	0	0	2	
I.2-26	N	CH	S	-	0	F	H	0	0	2	
I.2-27	CH	CH	S	-	0	CH ₃	CH ₃	0	0	2	1.75(s), 2.65 (m _c), 4.30(t), 7.35 (t), 7.45 (t), 7.75 (d), 7.90 (d)
I.2-28	CH	CH	S	-	0	(CH ₂) ₃ CH ₃	H	0	0	2	0.95(t), 1.30-1.55(m), 1.90-2.15 (m), 2.60(m _c), 4.30(t), 4.65(t), 7.30(t), 7.45 (t), 7.75 (d), 7.85 (d)
I.2-29	CH	CH	S	6-OCH ₂ CH ₃	1	(CH ₂) ₃ CH ₃	H	0	0	2	0.90(t), 1.30-1.50(m), 1.85-2.10 (m), 2.60(m _c), 4.05(q), 4.30(t), 4.50(t); 6.95(d), 7.20(s), 7.75 (d)

No.	B	D	E	(R ^c) _q	q	R ¹	R ²	m	n	p	Physical data: ¹ H-NMR (δ [ppm])
I.2-30	CH	CH	S	6-OCH ₂ CH ₃	1	CH ₃	CH ₃	0	0	2	1.45(t), 1.70(s), 2.60(m _c), 4.05(q), 4.30(t), 7.05(d), 7.25(s), 7.80(d)
I.2-31	CH	CH	S	6-OCH ₂ CH ₃	1	CH ₂ CH ₃	H	0	0	2	1.15(t), 1.45(t), 1.95-2.2(m), 4.05(q), 4.35(t), 4.5(t), 7.0(d), 7.2(s), 7.75(d)
I.2-32	N	CH	S	6-CF ₃	1	CH ₂ CH ₃	H	0	0	2	1.15(t), 2.0-2.25(m), 2.7(m _c), 4.35(q), 4.95(t), 8.4(s), 8.9(s)
I.2-33	CH	CH	O	-	0	CH ₃	H	0	0	2	1.75(d), 2.65(m _c), 4.35(t), 4.65(q), 7.25(m _c), 7.45(d), 7.6(d)
I.2-34	CH	CH	O	-	0	CH ₂ CH ₃	CH ₃	0	0	2	1.15(t), 2.10(m _c), 2.65(m _c), 4.3(t), 4.55(t), 7.25(m _c), 7.45(d), 7.60(d)
I.2-35	CH	CH	O	-	0	CH ₃	CH ₃	0	0	2	1.75(s), 2.65(m _c), 4.30(t), 7.25(m _c), 7.45(d), 7.60(d)
I.2-36	CH	CH	O	-	0	(CH ₂) ₃ CH ₃	H	0	0	2	0.95(t), 1.30-2.15(m), 2.65(m _c), 4.35(t), 4.55(t), 7.25(m _c), 7.45(d), 7.60(d)
I.2-37	CH	CH	O	-	0	CH(CH ₃) ₂	H	0	0	2	1.15(d), 2.40(m _c), 2.65(m _c), 4.35(t), 4.55(d), 7.25(m _c), 7.45(d), 7.60(d)
I.2-38	CH	N	S	-	0	CH ₃	H	0	0	2	1.75(d), 2.65(m _c), 4.35(t), 4.75(q), 7.35(m _c), 8.05(d), 8.45(d)
I.2-39	CH	N	S	-	0	CH ₃	CH ₃	0	0	2	1.75(s), 2.65(m _c), 4.30(t), 7.35(m _c), 8.05(d), 8.50(d)


No.	B	D	E	(R ^c) _q	q	R ¹	R ²	m	n	p	Physical data: ¹ H-NMR (δ [ppm])
I.2-40	CH	N	S	-	0	(CH ₂) ₃ CH ₃	H	0	0	2	0.95(t), 1.25-2.15(m), 2.65(m _c), 4.35(t), 4.75(t), 7.35(m _c), 8.0(d), 8.5(d)
I.2-41	N	CH	S	6-Cl	0	(CH ₂) ₃ CH ₃	H	0	0	2	0.95(t), 1.25-2.15(m), 2.65(m _c), 4.3(m _c), 4.95(t), 8.1(s), 8.55(s)
I.2-42	N	CH	S	-	0	CH(CH ₃) ₂	H	0	0	2	1.15(d), 2.4(m _c), 2.65(m _c), 4.35(m _c), 4.75(d), 7.35(m _c), 8.05(d), 8.45(d)
I.2-43	CH	CH	O	6-Cl	1	CH ₂ CH ₃	H	0	0	2	1.1(t), 2.05(m _c), 2.65(m _c), 4.35(t), 4.5(t), 7.25(d), 7.55(m _c)
I.2-44	CH	CH	O	6-Cl	1	CH ₃	CH ₃	0	0	2	1.80(s), 2.65(m _c), 4.3(t), 7.25(d), 7.45(s), 7.55(d)
I.2-45	CH	CH	O	6-Cl	1	(CH ₂) ₃ CH ₃	H	0	0	2	0.95(t), 1.25-2.15(m), 2.65(m _c), 4.3(t), 4.5(t), 7.25(d), 7.5(m _c)
I.2-46	CH	CH	S	6-OCH ₂ CH ₃	1	CH(CH ₃) ₂	H	0	0	2	1.15(m _c), 1.45(t), 2.35(m _c), 2.65(m _c), 4.05(q), 4.3(m _c), 4.5(d), 7.0(d), 7.2(s), 7.75(d)
I.2-47	CH	CH	S	6-CF ₃	1	H	H	0	0	2	
I.2-48	CH	CH	O	5-CH ₃	1	H	H	0	0	2	
I.2-49	CH	CH	S	6-F	1	H	H	0	0	2	
I.2-50	N	CH	S	6-CH ₃	1	F	H	0	0	2	
I.2-51	N	CH	S	6-CF ₃	1	CH ₃	H	0	0	2	
I.2-52	CH	CH	S	6-OCF ₃	1	H	H	0	0	2	

No.	B	D	E	(R ^o) _q		R ¹	R ²	m	n	p	Physical data: ¹ H-NMR (δ [ppm])
					q						
I.2-53	N	CH	S	6-Br	1	CH ₃	H	0	0	2	1.75(d), 2.65(m _c), 4.35(m _c), 4.95(q), 8.25(s), 8.65(s)
I.2-54	N	CH	S	6-Br	1	CH ₂ CH ₃	H	0	0	2	1.1(t), 2.10(m _c), 2.65(m _c), 4.35(m _c), 4.90(t), 8.25(s), 8.65(s)
I.2-55	N	CH	S	6-Br	1	CH ₃	CH ₃	0	0	2	1.85(s), 2.65(m _c), 4.35(t), 8.25(s), 8.65(s)
I.2-56	CH	CH	O	5-CH ₃	1	CH ₃	H	0	0	2	1.75(d), 2.45(s), 2.65(m _c), 4.35(t), 4.6(q), 7.05(d), 7.30(d), 7.4(s)
I.2-57	CH	CH	O	5-CH ₃	1	CH ₂ CH ₃	H	0	0	2	1.15(t), 2.1(m _c), 2.45(s), 2.65(m _c), 4.35(m _c), 4.35(t), 4.55(t), 7.05(d), 7.3(d), 7.4(s)
I.2-58	CH	CH	O	5-CH ₃	1	CH ₃	CH ₃	0	0	2	1.8(s), 2.45(s), 2.65(m _c), 4.35(m _c), 7.1(d), 7.3(d), 7.4(s)
I.2-59	CH	CH	S	5-Cl	1	CH(CH ₃) ₂	H	0	0	2	1.15(d), 2.4(m _c), 2.65(m _c), 4.35(m _c), 4.65(d), 7.25(d) 7.65(d), 7.85(d)
I.2-60	N	CH	S	-	0	CH ₂ CH ₃	H	0	0	2	1.1(t), 2.1(m _c), 2.7(m _c), 4.95(t), 7.25(m _c), 8.1(d), 8.65(d)
I.2-61	N	CH	S	-	0	CH ₃	CH ₃	0	0	2	1.85(s), 2.65(m _c), 4.35(m _c), 7.25(m _c), 8.1(d), 8.65(d)
I.2-62	N	CH	S	-	0	(CH ₂) ₃ CH ₃	H	0	0	2	0.95(t), 1.3-2.2(m), 2.65(m _c), 4.35(m _c), 4.95(t), 7.25(m _c), 8.1(d), 8.65(d)

No.	B	D	E	(R ^c) _q	q	R ¹	R ²	m	n	p	Physical data: ¹ H-NMR (δ [ppm])
I.2-63	CH	CH	S	5-Cl	1	CH ₃	H	0	0	2	1.75(d), 2.65(m _c), 4.3(t), 4.7(q), 7.25(d), 7.7(d), 7.85(s)
I.2-64	CH	CH	S	5-Cl	1	CH ₂ CH ₃	H	0	0	2	see example 3
I.2-65	CH	CH	S	5-Cl	1	CH ₃	CH ₃	0	0	2	1.75(s), 2.65(m _c), 4.35(t), 7.3(d), 7.7(d), 7.85(s)
I.2-66	CH	CH	S	5-Cl	1	(CH ₂) ₃ CH ₃	H	0	0	2	0.95(t), 1.3-2.1(m), 2.65(m _c), 4.35(m _c), 4.65(t), 7.3(d), 7.7(d), 7.85(s)
I.2-67	CH	CH	S	5-OCH ₃	1	H	H	0	0	2	2.85(m _c), 3.55(t), 3.9(s), 6.95(d), 7.4(s), 7.6(d)
I.2-68	CH	CH	S	5-OCH ₃	1	CH ₃	H	0	0	2	1.7(d), 2.65(m _c), 3.85(s), 4.35(m _c), 4.7(q), 6.95(d), 7.4(s), 7.6(d)
I.2-69	CH	CH	S	5-OCH ₃	1	CH(CH ₃) ₂	H	0	0	2	1.15(m _c), 2.4(m _c), 2.65(m _c), 3.85(s), 4.35(t), 4.6(d), 6.95(d), 7.4(s), 7.6(d)
I.2-70	CH	N	NCH ₃	-	0	H	H	0	0	2	2.85(m _c), 3.65(t), 3.75(s), 7.2(m _c), 7.85(d), 8.25(d)
I.2-71	CH	CH	O	6-Cl	1	H	H	0	1	2	2.65(m _c), 3.0(t), 3.55(m _c), 4.35(t), 7.25(d), 7.45(s), 7.5(d)
I.2-72	CH	CH	S	-	0	H	H	0	2	2	
I.2-73	CH	CH	S	6-OCH ₂ CH ₃	1	H	H	0	2	2	1.45(t), 2.15(m _c), 2.5(t), 2.65(m _c), 3.35(t), 4.05(q), 4.25(t), 7.0(d), 7.2(s), 7.75(d)

No.	B	D	E	(R ^c) _q	q	R ¹	R ²	m	n	p	Physical data: ¹ H-NMR (δ [ppm])
I.2-74	CH	CH	O	-	0	H	H	0	2	2	2.2(m _c), 2.5-2.7(m), 3.35(m _c), 4.25(m _c), 7.25(m _c), 7.45(d), 7.6(d)
I.2-75	CH	CH	O	6-Cl	1	H	H	0	2	2	2.2(m _c), 2.55(m _c), 2.65(m _c), 3.35(m _c), 4.3(m _c), 7.25(d), 7.45(s), 7.5(d)
I.2-76	CH	N	S	-	0	CH ₃	H	2	0	2	1.8(d), 2.55(m _c), 4.3(m _c), 4.65(q), 7.65(m _c), 8.5(d), 8.8(d)
I.2-77	CH	N	S	-	0	CH ₃	CH ₃	2	0	2	1.8(s), 2.65(m _c), 4.3(t), 7.6(dd), 8.5(d), 8.8(d)
I.2-78	CH	N	S	-	0	CH ₃	CH ₃	1	0	2	1.85(s), 2.65(m _c), 4.35(t), 7.55(dd), 8.15(d), 8.4(d)

Table II




No.	E'	(R ^c) _q	q	R ¹	R ²	m	n	p	Physical data: ¹ H-NMR (δ [ppm])
I.3-1	S	4-(p-F-C ₆ H ₄)	1	CH ₂ CH ₃	CH ₂ CH ₃	0	0	2	1.0(m _c), 2.0(m _c), 2.6(m _c), 4.3(m _c), 7.1(m _c), 7.4(s), 7.85(m _c)
I.3-2	S	4-(p-F-C ₆ H ₄)	1	H	H	0	0	2	2.65(m _c), 4.1(s), 4.35(m _c), 7.1(m _c), 7.3(s), 7.85(m _c)
I.3-3	S	4-(p-F-C ₆ H ₄)	1	CH ₃	H	0	0	2	1.7(d), 2.6(m _c), 4.3(m _c), 4.45(q), 7.1(m _c), 7.35(s), 7.85(m _c)

No.	E'	(R ^c) _q	q	R ¹	R ²	m	n	p	Physical data: ¹ H-NMR (δ [ppm])
I.3-4	S	4-(p-F-C ₆ H ₄)	1	CH ₃	CH ₃	0	0	2	1.7(s), 2.6(m _c), 4.25(t), 7.1(m _c), 7.4(s), 7.85(m _c)
I.3-5	S	4-(p-F-C ₆ H ₄)	1	CH(CH ₃) ₂	H	0	0	2	1.15(m _c), 2.3(m _c), 2.65(m _c), 4.2-4.4(m), 7.1(m _c), 7.3(s), 7.8(m _c)
I.3-6	S	4-(p-F-C ₆ H ₄)	1	(CH ₂) ₃ CH ₃	H	0	0	2	1.95(m _c), 1.2-2.1(m), 2.65(m _c), 4.25-4.4(m), 7.1(m _c), 7.35(s), 7.85(m _c)
I.3-7	S	4-(p-F-C ₆ H ₄)	1	CH ₂ CH ₃	H	0	0	2	1.15(t), 2.05(m _c), 2.6(m _c), 4.2-4.45(m), 7.1(t), 7.3(s), 7.85(m _c)
I.3-8	S	4-(p-Cl-C ₆ H ₄)	1	CH ₃	CH ₃	0	0	2	1.7(s), 2.55(m _c), 4.25(t), 7.4(d), 7.5(s), 7.8(d)
I.3-9	S	4-(p-Cl-C ₆ H ₄)	1	CH ₃	H	0	0	2	1.7(s), 2.6(m _c), 4.3(m _c), 4.45(q), 7.35(m _c), 7.8(d)
I.3-10	S	4-(p-Cl-C ₆ H ₄)	1	(CH ₂) ₃ CH ₃	H	0	0	2	see example 4
I.3-11	S	4-(p-Cl-C ₆ H ₄)	1	CH ₂ CH ₃	H	0	0	2	1.15(t), 2.05(m _c), 2.65(m _c), 4.25-4.4(m), 7.35(m _c), 7.8(d)
I.3-12	S	4-(o,p-Cl ₂ -C ₆ H ₃)	1	CH ₃	H	0	0	2	1.65(m _c), 2.6(m _c), 4.2-4.5(m), 7.35(d), 7.5(s), 7.8(c), 7.95(d)
I.3-13	S	4-(o,p-Cl ₂ -C ₆ H ₃)	1	CH ₃	CH ₃	0	0	2	1.7(s), 2.6(m _c), 4.3(m _c), 7.35(d), 7.5(s), 7.9(m _c)
I.3-14	S	4-(o,p-Cl ₂ -C ₆ H ₃)	1	(CH ₂) ₃ CH ₃	H	0	0	2	0.9(t), 1.3-2.1(m), 2.6(m _c), 4.2-4.4(m), 7.3(d), 7.5(s), 7.8(s), 7.95(d)

No.	E'	(R ^c) _q	q	R ¹	R ²	m	n	p	Physical data: ¹ H-NMR (δ [ppm])
I.3-15	S	4-(o,p-Cl ₂ -C ₆ H ₃)	1	CH(CH ₃) ₂	H	0	0	2	1.1(dd), 2.3(m _c), 2.65(m _c), 4.2-4.4(m), 7.3(d), 7.5(s), 7.8(s), 7.95(d)
I.3-16	S	4-(p-Cl-C ₆ H ₄)	1	CH(CH ₃) ₂	H	0	0	2	1.15(dd), 2.35(m _c), 2.65(m _c), 4.2-4.4(m), 7.35(m _c), 7.8(d)
I.3-17	S	4-(C ₆ H ₅)	1	CH ₃	CH ₃	0	0	2	1.7(s), 2.6(m _c), 4.3(t), 7.3-7.5(m), 7.85(d)
I.3-18	S	4-(C ₆ H ₅)	1	(CH ₂) ₃ CH ₃	H	0	0	2	0.9(t), 1.3-2.1(m), 2.6(m _c), 4.2-4.45(m), 7.25-7.45(m), 7.85(d)
I.3-19	S	5-(C ₆ H ₅)	1	CH(CH ₃) ₂	H	0	0	2	1.15(dd), 2.35(m _c), 2.65(m _c), 4.2-4.4(m), 7.25-7.45(m), 7.85(d)
I.3-20	O	4-(C ₆ H ₅), 5-(C ₆ H ₅)	2	H	H	0	0	2	2.65(m _c), 4.05(s), 4.35(t), 7.25-7.60(m)
I.3-21	O	4-(C ₆ H ₅), 5-(C ₆ H ₅)	2	CH ₂ CH ₃	H	0	0	2	1.15(t), 2.1(m _c), 2.65(m _c), 4.3(m _c), 7.25-7.6(m)
I.3-22	O	4-(C ₆ H ₅), 5-(C ₆ H ₅)	2	CH ₃	H	0	0	2	1.75(d), 2.65(m _c), 4.25-4.5(m), 7.25-7.55(m)
I.3-23	O	4-(C ₆ H ₅), 5-(C ₆ H ₅)	2	CH ₃	CH ₃	0	0	2	1.75(s), 2.65(m _c), 4.3(t), 7.25-7.6(m)
I.3-24	O	4-(C ₆ H ₅), 5-(C ₆ H ₅)	2	CH(CH ₃) ₂	H	0	0	2	1.15(dd), 2.35(m _c), 2.65(m _c), 4.25-4.45(m), 7.25-7.6(m)
I.3-25	O	4-(C ₆ H ₅), 5-(C ₆ H ₅)	2	(CH ₂) ₃ CH ₃	H	0	0	2	0.9(t), 1.3-2.1(m), 2.65(m _c), 4.25-4.45(m), 7.25-7.6(m)
I.3-26	S	4-(C ₆ H ₅)	1	H	H	0	0	2	

No.	E'	(R ^c) _q	q	R ¹	R ²	m	n	p	Physical data: ¹ H-NMR (δ [ppm])
I.3-27	S	4-(C ₆ H ₅)	1	CH ₃	H	0	0	2	
I.3-28	S	4-(C ₆ H ₅)	1	CH ₂ CH ₃	H	0	0	2	
I.3-29	S	4-(C ₆ H ₅)	1	H	H	0	1	2	2.65(m _c), 2.95(t), 3.55(t), 4.35(t), 7.3-7.55(m), 7.85(d)
I.3-30	S	4-(C ₆ H ₅)	1	H	H	0	2	2	2.15(m _c), 2.5-2.7(m), 3.35(t), 4.25(t), 7.25-7.45(m), 7.85(d)
I.3-31	S	4-(p-F-C ₆ H ₄)	1	H	H	2	0	2	2.65(m _c), 4.3(t), 4.55(s), 7.15(t), 7.8(s), 7.95(m _c)
I.3-32	S	4-(p-F-C ₆ H ₄)	1	CH ₃	H	2	0	2	1.8(d), 2.65(m _c), 4.3(m _c), 4.55 (q), 7.15(m _c), 7.8(s), 7.95(m _c)
I.3-33	S	4-(p-F-C ₆ H ₄)	1	CH ₂ CH ₃	H	1	0	2	1.1(m _c), 2.15(m _c), 2.6(m _c), 3.85-4.4(m), 7.1(m _c), 7.75(d), 7.85(m _c)
I.3-34	S	4-(p-F-C ₆ H ₄)	1	CH ₃	CH ₃	1	0	2	1.2(s), 1.7(s), 2.65(m _c), 4.35(m _c), 7.1(m _c), 7.7(s), 7.8(m _c)
I.3-35	S	4-(p-F-C ₆ H ₄)	1	(CH ₂) ₃ CH ₃	H	1	0	2	0.9(m _c), 1.15-2.2(m), 2.6(m _c), 3.9-4.4(m), 7.15(m _c), 7.7(d), 7.85(m _c)

Table III



No.	E	R ^c	R ¹	R ²	p	Physical data: ¹ H-NMR (δ [ppm])
I.4-1	O	p-F-C ₆ H ₄	H	H	2	2.7 (m _c), 4.15 (s), 4.4 (t), 7.2 (m _c), 8.0 (m _c)
I.4-2	O	p-CH ₃ -C ₆ H ₄	H	H	2	2.4 (s), 2.65 (m _c), 4.15 (s), 4.35 (t), 7.3 (m _c), 7.9 (m _c)
I.4-3	O	p-F-C ₆ H ₄	CH ₃	H	2	1.75 (d), 2.65 (m _c), 4.25-4.55 (m), 7.2 (m _c), 8.0 (m _c)
I.4-4	O	p-F-C ₆ H ₄	CH(CH ₃) ₂	H	2	1.15 (m _c), 2.4 (m _c), 2.65 (m _c), 4.3-4.45 (m), 7.2 (m _c), 8.0 (m _c)
I.4-5	S	CH ₃	H	H	2	2.65 (m _c), 2.75 (s), 4.15 (s), 4.35 (t)
I.4-6	S	CH ₃	CH ₃	H	2	1.7 (d), 2.65 (m _c), 2.75 (s), 4.35 (m _c), 4.55 (q)
I.4-7	S	CH ₃	CH(CH ₃) ₂	H	2	1.15 (dd), 2.35 (m _c), 2.65 (m _c), 2.75 (s), 4.3 (m _c), 4.45 (d)
I.4-8	O	p-F-C ₆ H ₄	CH ₂ CH ₃	H	2	1.15 (t), 2.15 (m _c), 2.65 (m _c), 4.3-4.5 (m), 7.25 (m _c), 8.0 (m _c)
I.4-9	S	CH ₃	CH ₂ CH ₃	H	2	1.1 (t), 2.05 (m _c), 2.65 (m _c), 2.75 (s), 4.25-4.5 (m)
I.4-10	S	CH ₃	CH ₃	H	2	1.75 (d), 2.65 (m _c), 4.35 (m _c), 4.75 (q), 9.05 (s)
I.4-11	S	CH ₃	H	H	2	2.65 (m _c), 4.2 (s), 4.35 (m _c), 9.05 (s)

Examples of action against animal pests

The action of the compounds of the formula I against pests was
5 demonstrated by the following experiments:

The active compounds were formulated

- a. for testing the activity against nematodes as an about 0.05%
w/v strength solution in a carrier of 5% by volume of acetone
10 and 0.05% by volume Tween 20 (Polyoxyethylene-(20)-sorbitan
monolaurate) in water, which was then diluted with 5% acetone
in water and 0.05% Tween 20 in water to achieve the desired
concentrations for drenching.
- b. for testing the activity against insects and arachnids as a
15 10.000 ppm solution in a mixture of 35% acetone and water,
which was diluted with water, if needed.

After the experiments were completed, in each case the lowest
concentration was determined at which the compound still caused
20 an 80 to 100% inhibition or mortality in comparison with
untreated controls (limit or minimal concentration).

Activity against nematode plant diseases

- 25 Soybean cyst nematode (SCN), *Heterodera glycines*
Tomato root knot nematode (RKN), *Meloidogyne incognita*

Silty loam soil in individual pots with 1-week-old tomato
transplants (cultivar Bonny Best) and soybean transplants
30 (cultivar Hutcheson) were drenched with the test solution of the
active ingredient. Aqueous suspensions of J2 nematode larvae and
Meloidogyne incognita in the case of tomatoes and *Heterodera*
glycines in the case of soybeans were drenched on the soil
surface. Plants were kept one day in a moist infection chamber at
35 26°C then moved into the greenhouse and maintained with bottom
watering until harvested for evaluation. The tests were
replicated 3 times for each disease.

In the case of root knot nematode on tomatoes, plants were
40 harvested two weeks after treatment and inoculation. Roots were
washed free of soil and the number of root knot galls on each
root system was visually examined and compared for treated and
plants that were only treated with the acetone / Tween 20
carrier.

114

In the case of soybean cyst nematode on soybeans, plants were harvested four weeks after treatment and inoculation. Roots were washed free of soil and the number of cysts per root-mass was visually examined and compared for treated and plants that were
5 only treated with the acetone / Tween 20 carrier.

In this test, tomato plants which had been treated with 0.6 kg/ha of the compounds I.2-11, I.2-29, I.2-31, I.2.-32, I.2-33, I.2-37, I.3.-2, I.3.-4, I.3-7 and I.3-11 showed a reduction in root knot
10 galling of 100% compared to plants treated with the acetone/Tween 20 carrier.

Activity against insects and arachnids

15 *Spodoptera eridania*, 2nd instar larvae, southern armyworm (SAW)

Leaves of two lima bean plants contained in pots at BBCH stage 11 were dipped in the test solution, allowed to dry and then placed in plastic bags with holes punched for ventilation. Ten 2nd
20 instar larvae were introduced. After 4 days, mortality, reduced feeding, or any interference with normal growth was examined visually.

Diabrotica virgifera virgifera Leconte, 2nd instar western corn
25 rootworm (WCR)

1 ml of the test solution was pipetted onto 1 ccm³ of talc in a 30 ml screw-top glass jar so as to provide 1.25 mg of active ingredient per jar (corresponding to about 50 kg/ha). The dried
30 talc was loosened, and 1 ccm³ of millet seed as food for the insects and 25 ml of moist soil were added to each jar and the contents were mixed mechanically. 10 2nd instar rootworms were added to each jar and the jars are loosely capped to allow air exchange for the larvae. The treatments were held for 5 days
35 when mortality counts were made. Missing larvae were presumed dead, since they decompose rapidly and cannot be found.

Tetranychus urticae (OP-resistant strain), 2-spotted spider mite (TSM)
40

Sieva lima bean plants with primary leaves expanded to 7-8 cm were infested by placing on each a small piece from an infested leaf (with about 100 mites) taken from the main colony. This was done at about 2 hours before treatment to allow the mites to move
45 over to the test plant to lay eggs. The piece of leaf used to transfer the mites was removed. The newly-infested plants were

115

dipped in the test solution and allowed to dry. After 2 days, one leaf is removed and mortality counts are made.

Aphis gossypii, cotton aphid (CA)

5

Cotton plants at the cotyledon stage (one plant per pot) were infested by placing a heavily infested leaf from the main colony on top of each cotyledon. The aphids were allowed to transfer to the host plant overnight, and the leaf used to transfer the

10 aphids was removed. The cotyledons were dipped in the test solution and allowed to dry. After 5 days, mortality counts were made.

Spodoptera eridania, eggs-southern armyworm and *Diabrotica*

15 *undecimpunctata howardi*, eggs-southern corn rootworm (SAW-Eggs) and (SCR-Eggs)

Wells containing artificial diet were treated with the test solutions and dried. The appropriate insect eggs were then placed
20 in the wells which were covered with vented covers. After 7 days, mortality counts were made.

In this test, *Tetranychus urticae* which had been treated with 100 ppm of compound I.2-6 showed a kill rate of over 75% whereas

25 untreated pests showed a rate of 0%.

30

35

40

45